

CHAPTER II.

PHYSIOGRAPHY.

§ 1. General Description of Australia.

1. Geographical Position.—(i) *General.* The Australian Commonwealth, which includes the island continent of Australia proper and the island of Tasmania, is situated in the Southern Hemisphere, and comprises in all an area of about 2,974,581 square miles, the mainland alone containing about 2,948,366 square miles. Bounded on the west and east by the Indian and Pacific Oceans respectively, it lies between longitudes 113° 9' E. and 153° 39' E., while its northern and southern limits are the parallels of latitude 10° 41' S. and 39° 8' S., or, including Tasmania, 43° 39' S. On its north are the Timor and Arafura Seas and Torres Strait—on its south the Southern Ocean and Bass Strait. The extreme points are “Steep Point” on the west, “Cape Byron” on the east, “Cape York” on the north, “Wilson’s Promontory” on the south, or, if Tasmania be included, “South-East Cape.”

(ii) *Tropical and Temperate Regions.* Of the total area of Australia nearly 40 per cent. lies within the tropics. Assuming, as is usual, that the latitude of the Tropic of Capricorn is 23° 30' S., the areas within the tropical and temperate zones are approximately as follows :—

AUSTRALIA : AREAS OF TROPICAL AND TEMPERATE REGIONS.

Area.	N.S.W. (a)	Vic.	Qld.	S. Aust.	W. Aust.	Tas.	N. Terr.	Total.
Within Tropical Zone sq. miles	359,000	..	364,000	..	426,320	1,149,320
Within Temperate Zone sq. miles	310,372	87,884	311,500	380,070	611,920	26,215	97,300	1,825,261
Total Area sq. miles	310,372	87,884	670,500	380,070	975,920	26,215	523,620	2,974,581
Ratio of Tropical part whole State	0.535	..	0.373	..	0.814	0.386
Ratio of Temperate part to whole State	1	1	0.465	1	0.627	1	0.186	0.614

(a) Includes Australian Capital Territory.

Thus, of the whole of Australia, the tropical part is roughly about five-thirteenths, (0.386) or, of the three territories with areas within the tropical zone, about one-half (0.530).

2. Area of Australia compared with Areas of other Countries.—The area of Australia is almost as great as that of the United States of America, four-fifths of that of Canada, more than one-fifth of the area of the British Commonwealth, nearly three-fourths of the whole area of Europe, and about 25 times as large as Great Britain and

Ireland. The areas of Australia and of certain other countries are shown in the following table :—

AREA OF AUSTRALIA AND OF OTHER COUNTRIES, *circa* 1949.
(’000 sq. miles.)

Country.	Area.	Country.	Area.
Continental Divisions—		Africa—<i>continued</i>.	
Europe (<i>a</i>)	1,914	Angola	481
Asia (<i>a</i>)	10,359	Union of South Africa	472
U.S.S.R. (Europe and Asia)	8,599	Ethiopia	409
Africa	11,681	Egypt	386
North and Central America and West Indies	9,367	Tanganyika Territory	363
South America	6,854	Nigeria and Protectorate.. ..	339
Oceania	3,304	South-West Africa	318
Total, excluding Arctic and Antarctic Conts.	52,078	Mozambique	298
		Northern Rhodesia	290
		Bechuanaland Protectorate	275
		Madagascar	229
Europe(<i>a</i>)—		Kenya Colony and Protec- torate	225
France	213	Other	1,420
Spain (incl. possessions)	194	Total	11,681
Sweden	173		
Germany	136	North and Central America—	
Finland	130	Canada	3,843
Norway	125	United States of America.. ..	3,022
Poland	120	Greenland	840
Italy	116	Mexico	760
Yugoslavia	99	Alaska	586
United Kingdom	94	Honduras	59
Rumania	92	Nicaragua	57
Other	422	Other	200
Total	1,914	Total	9,367
Asia(<i>a</i>)—		South America—	
China and Dependencies	3,759	Brazil	3,288
India	1,221	Argentine Republic	1,079
Indonesia(<i>b</i>)	735	Peru	482
Iran	629	Colombia (excl. of Panama)	440
Mongolian Peoples' Republic	626	Bolivia	413
Saudi Arabia	597	Venezuela	352
Pakistan	360	Chile	286
Turkey	287	Paraguay	157
French Indo-China	272	Ecuador	106
Burma	262	Other	251
Afghanistan	251	Total	6,854
Thailand	198		
Other	1,162	Oceania—	
Total	10,359	Commonwealth of Australia	2,975
U.S.S.R.	8,599	New Zealand and Depen- dencies	104
Africa—		New Guinea	93
French West Africa	1,805	Papua	91
French Equatorial Africa	969	Other	41
Anglo-Egyptian Sudan	967	Total	3,304
Belgian Congo	905		
Algeria	851	British Commonwealth	13,258
Libya	679		

(a) Excludes U.S.S.R., shown below.

(b) Includes Dutch New Guinea.

The areas shown in the table are obtained from the *Demographic Yearbook*, 1949-50 published by the United Nations and the countries have been arranged in accordance with the continental groups used therein.

3. Areas of Political Subdivisions.—As already stated, Australia consists of six States and the Northern, and Australian Capital, Territories. The areas of these, and their proportions of the total of Australia, are shown in the following table :—

AUSTRALIA : AREA OF STATES AND TERRITORIES.

State or Territory.	Area.	Proportion of Total.
	Sq. miles.	%
New South Wales	309,433	10.40
Victoria	87,884	2.96
Queensland	670,500	22.54
South Australia	380,070	12.78
Western Australia	975,920	32.81
Tasmania	26,215	0.88
Northern Territory	523,620	17.60
Australian Capital Territory	939	0.03
Total	2,974,581	100.00

4. Coastal Configuration—(i) *General*. There are no striking features in the configuration of the coast ; the most remarkable indentations are the Gulf of Carpentaria on the north and the Great Australian Bight on the south. The Cape York Peninsula on the extreme north is the only other remarkable feature in the outline. In Official Year Book No. 1 an enumeration is given of the features of the coast-line of Australia (see pp. 60-68).

(ii) *Coast-line*. The lengths of coast-line, excluding minor indentations, of each State and of the whole continent, and the area per mile of coast-line, are shown in the following table :—

AUSTRALIA : COAST-LINE AND AREA PER MILE THEREOF.

State.	Coast-line.	Area per Mile of Coast-line.	State.	Coast-line.	Area per Mile of Coast-line
	Miles.	Sq. miles.		Miles.	Sq. miles.
New South Wales(a)	700	443	Western Australia	4,350	224
Victoria	680	129	Northern Territory	1,040	503
Queensland	3,000	223	Continent (b) ..	11,310	261
South Australia	1,540	247	Tasmania	900	29

(a) Includes Australian Capital Territory.

(b) Area 2,948,366 square miles.

For the entire Commonwealth of Australia there is a coast-line of 12,210 miles with an average of 244 square miles for one mile of coast-line. According to Strelbitaki, Europe has only 75 square miles of area to each mile of coast-line, and, according to more recent figures, England and Wales have only one-third of this, 25 square miles.

5. Geographical Features of Australia.—In separate issues of earlier Official Year Books fairly complete information has been given concerning some special geographical element. The nature of this information and its position in the various issues can be readily ascertained on reference to the special index following the index to maps and graphs at the end of this issue.

6. Fauna, Flora, Geology and Seismology of Australia.—Special articles dealing with these features have appeared in previous issues of the Official Year Book, but limits of space naturally preclude their repetition in each volume. As pointed out in par. 5, however, the nature and position of these articles can be readily ascertained from the special index.

§ 2. Climate and Meteorology of Australia.*

1. **Introductory.**—Previous issues of the Official Year Book, notably No. 3, pp. 79 and 80, and No. 4, pp. 84 and 87, contained outlines of the history of Australian meteorology and the creation and organization of the Commonwealth Bureau of Meteorology. Official Year Book No. 38, pp. 30-32, contained paragraphs devoted to (i) Organization of the Meteorological Service; (ii) Meteorological Publications; (iii) Equipment; and (iv) Meteorological Divisions.

By reason of its insular geographical position and the absence of striking physical features, whether in marine gulfs or in important mountains, Australia is, on the whole, less subject to extremes of weather than are regions of similar area in other parts of the globe, and latitude for latitude Australia is, on the whole, more temperate.

The average elevation of the surface of the land is low, probably close to 900 feet above the sea. The altitudes range up to a little over 7,300 feet, hence its climate embraces a great many features, from the characteristically tropical to what is essentially alpine, a fact indicated in some measure by the name Australian Alps given to the southern portion of the Great Dividing Range.

On the coast, the rainfall is often abundant and the atmosphere moist, but in some portions of the interior is very limited, and the atmosphere dry. The distribution of forest, therefore, with its climatic influence, is very uneven. In the interior, in places, there are fine belts of trees, but there are large areas also which are treeless, and here the air is hot and parching in summer. Again, on the coast, even so far south as latitude 35°, the vegetation is tropical in its luxuriance, and to some extent also in character. Climatologically, therefore, Australia may be said to present a great variety of features.

2. **Temperature.**—(i) *Effective Temperature.* When a meteorologist speaks of temperature he means the temperature of the air indicated by a thermometer sheltered from precipitation, from direct rays of the sun and from radiation of heat from the ground and neighbouring objects, yet freely exposed to the circulation of the air. In other words, he means temperature measured under conditions standardized as near as possible in a Stevenson Screen, which is the standard housing for meteorological thermometers.

This shade temperature as measured by a "dry bulb" thermometer shows only the actual temperature experienced by dry inorganic substances, not the *sensible* temperatures felt by organic bodies. In the case of human beings, sensible temperature is affected by the rate of conduction of heat to or from the body by moving air and also by the rate of cooling due to evaporation from the skin and respiratory passages. The wind and humidity therefore determine the sensible temperature.

The humidity (relative humidity) is determined from the readings of the dry and wet bulb thermometers. Of late years, however, with increasing interest in human comfort in tropical climates, another term, *effective* temperature, has come into use. It may be defined as "the temperature of a still, saturated atmosphere which would on the average produce the same feeling of warmth or cold as the atmosphere in question".†

Later investigations have established "comfort zones"‡ bounded by limits of effective temperature within which people will feel comfortable. American research workers have determined the following figures :—§

COMFORT ZONES : EFFECTIVE TEMPERATURES.

Season.	No subjects feel comfortable below—	Fifty per cent. of subjects feel comfortable between—	No subjects feel comfortable above—
Winter	60° F.	63° and 71° F.	74° F.
Summer	64° F.	66° and 75° F.	79° F.

* Prepared from data supplied by the Acting Director, Commonwealth Meteorological Bureau.
 † Houghton, F. C., Teague, W. W. and Miller, W. E. (1926) Amer. Soc. Heat. Vent. Engns.
 ‡ Yaglou, C. P. (1926) J. Industr. Hyg. § Yaglou, C. P. (1927) Ibid.

Isotherms of effective temperature (not corrected for altitude) have been determined for Australia.* A map showing effective temperature for Australia for January (9 a.m.) will be found on page 33.

It will be seen that the 80° F. isotherm is confined to a very narrow tract of country on the north-west coast of Western Australia. The 75° F. isotherm extends broadly from Onslow on the north-west coast of Western Australia to Daly Waters to Camoowal to Moreton in Cape York Peninsula following in a general way the coastline of Northern Australia but from 100 to 300 miles inland.

Queensland investigators† in recent years have divided some towns of Queensland into three classes on the basis of deviation from comfort:—

Class 1 (Sub-tropics).—Quite suitable for Caucasian habitation—Rockhampton, Bundaberg, Brisbane, Longreach, Charleville.

Class 2 (Marginal tropics).—Suitable for Caucasian habitation, but requires adaptation in summer—Mackay, Townsville.

Class 3 (Tropics).—(a) Permissible for Caucasian habitation but requires selection and marked adaptation—Cardwell, Cairns, Cloncurry. (b) Not suitable for continuous Caucasian habitation—Cape York, Burketown.

These results of recent years bear out investigations made previously in Australia‡ in which the atmospheric vapour pressure was used as a measure of comfort, its value for this purpose being that it has equal effect in both indoor and outdoor climates. The limits of comfort range from .2 to .5 inch of vapour pressure. After drawing isopleths for effective temperature (not corrected for altitude), mean vapour pressure reduced to a logarithmic scale, and mean wet bulb, it is found that there is close agreement in defining zones of relative discomfort.

(ii) *Seasons.* The Australian seasons are:—Summer, December to February; autumn, March to May; winter, June to August; spring, September to November. In most parts of Australia, January is the hottest month, but in Tasmania and southern Victoria, February is the hottest; in the tropical north, probably because the cooling "monsoon" rains occur in late summer, December is the hottest month, and at Darwin, November.

On a rainfall basis, in the tropical north the year is divisible into "wet" and "dry" seasons, but on the basis of temperature and physical comfort the "dry" season can be further sub-divided into two parts—"cool dry" and "warm dusty". §

(a) "*Cool dry*" Season. From May to August. The average maximum temperature ranges from 80° to 85° F., the relative humidity is low and in inland areas cold nights are experienced when the temperature drops to 40° F. The skies generally are cloudless, but in about one year in three during June or July one to two inches of rain fall.

(b) "*Warm dusty*" Season. From the end of August temperatures rise and reach a maximum in October or the beginning of November. Temperatures of over 120° F. have been recorded.

(c) "*Wet*" Season. After the first of the heavy storms, the maximum temperatures fall but still remain high with high relative humidity. At Wyndham during January, 1944 the minimum temperature did not drop below 75° F. for fourteen consecutive days. A maximum of over 100° F. was recorded on each rainless day.

In Central as in Northern Australia during the hottest months, the average temperatures range from 80° to 85° F., whereas in Southern Australia they vary from 65° to 70°.

Throughout Australia the coldest month is July, when only a very narrow strip of the northern sea-board has an average temperature as high as 75°. Over the southern half of the continent, July temperatures range from 55° to 45° at elevations below 1,500

* Hounam, C. E. Effective Temp. Data, C.W.B. unpublished. † Lee, D. H. K. Trans. Roy. Soc. Trop. Med. and Hyg. (1940) Vol. XXXII.

‡ Barkley, H. Zones of Relative Physical Comfort in Australia. Met. Bull. 20, 1934. § Maze, W. H. Aust. Geog. June, 1945. Settlement in E. Kimberleys.

feet and fall as low as 35° on the Australian Alps. Here the temperature seldom, if ever, reaches 100° even in the hottest of seasons. Hotham Heights (6,100 feet above Mean Sea Level) recorded the highest maximum of 82.0° on 19th January, 1935. In winter, readings slightly below zero are occasionally recorded on the extreme heights.

Tasmania as a whole enjoys a moderate and equable range of temperature throughout the year, although occasionally hot winds may cause the temperature to rise to 100° in the eastern part of the State.

(iii) *Comparisons with other Countries.* In respect of Australian temperatures generally, it may be pointed out that the mean annual isotherm for 70° F. extends in South America and South Africa as far south as latitude 33°, while in Australia it reaches only as far south as latitude 30°, thus showing that, on the whole, Australia has, latitude for latitude, a more temperate climate than other places in the Southern Hemisphere.

The comparison is even more favourable when the Northern Hemisphere is included, for in the United States of America the 70° isotherm extends in several of the western States as far north as latitude 41°. In Europe, the same isotherm reaches almost to the southern shores of Spain, passing afterwards, however, along the northern shores of Africa till it reaches the Red Sea, when it bends northward along the eastern shore of the Mediterranean till it reaches Syria. In Asia, nearly the whole of the land area south of latitude 40° N. has a higher temperature than 70°.

The extreme range of temperature is less than 100° over practically the whole of Australia, that figure being only slightly exceeded at a very few places; it is mostly 70° to 90° over inland areas, and somewhat less on the coast. In parts of Asia and North America, the extreme range exceeds 130° and 150° in some localities.



Along the northern shores of Australia the temperatures are very equable. At Darwin, for example, the difference in the means for the hottest and coldest month is only 8.4°, and the extreme readings for the year, or the highest maximum on record and the lowest minimum, show a difference of under 50°.

The highest temperature recorded in Australia was 127.5° F. at Cloncurry on 16th January, 1889. The world's highest (136° F.) was recorded at Azizia (Tripoli) on 13th August, 1922. The lowest temperature ever recorded in Australia was -8° F. at Charlotte Pass on 14th June, 1945, and again on 22nd July, 1947, as contrasted with the world's lowest recorded temperature of -90° F. at Verkhoyansk (Siberia) on 5th and 7th February, 1892.


A comparison of the mean temperatures and the range from the extreme maximum to the extreme minimum temperatures (in whole degrees) of the capital cities of Australia with those of the main cities of some other countries is shown in tabular form in Official Year Book No. 38, p. 42.

(iv) *Hottest and Coldest Parts.* A comparison of the temperatures recorded at coast and inland stations shows that, in Australia, as in other continents, the range increases, within certain limits, with increasing distance from the coast. This is clearly illustrated by the map of extreme temperature range (page 33).


In the interior of Australia, and during exceptionally dry summers, the temperature occasionally reaches or exceeds 120° in the shade. The hottest area of the continent is situated in the northern part of Western Australia about the Marble Bar and Nullagine gold-fields, where the maximum shade temperature during the summer sometimes exceeds 100° continuously for days and weeks. The longest recorded period was 160 days from 31st October, 1923 to 7th April, 1924.




AREA AFFECTED AND PERIOD OF DURATION OF THE LONGEST HEAT WAVES WHEN THE MAXIMUM TEMPERATURE FOR CONSECUTIVE 24 HOURS REACHED OR EXCEEDED 100°F.




NOTE: THE FIGURES ON THE LINES INDICATE THE NUMBER OF CONSECUTIVE DAYS.





STATE LIBRARY OF VICTORIA



REFERENCE TO DISTRICTS

WESTERN AUSTRALIA	OVERLAND	NEW SOUTH WALES
1 North Kimberley	16 (Cooberdy)	21 Sydney
2 East Kimberley	17 Central Kimberley	22 Sydney
3 West Kimberley	18 Upper Western	23 Sydney
4 Gascoyne	19 Lower Western	24 Sydney
5 Pilbara	20 Port Curtis	25 Sydney
6 Great Southern	21 East Darling Downs	26 Sydney
7 South Coast	22 West Darling Downs	27 Sydney
8 Central South	23 Darling Downs	28 Sydney
9 Central West	24 Darling Downs	29 Sydney
10 Central North	25 Darling Downs	30 Sydney
11 North West	26 Darling Downs	31 Sydney
12 North East	27 Darling Downs	32 Sydney
13 Northern Plains	28 Darling Downs	33 Sydney
14 Northern Plains	29 Darling Downs	34 Sydney
15 Northern Plains	30 Darling Downs	35 Sydney
16 Northern Plains	31 Darling Downs	36 Sydney
17 Northern Plains	32 Darling Downs	37 Sydney
18 Northern Plains	33 Darling Downs	38 Sydney
19 Northern Plains	34 Darling Downs	39 Sydney
20 Northern Plains	35 Darling Downs	40 Sydney
21 Northern Plains	36 Darling Downs	41 Sydney
22 Northern Plains	37 Darling Downs	42 Sydney
23 Northern Plains	38 Darling Downs	43 Sydney
24 Northern Plains	39 Darling Downs	44 Sydney
25 Northern Plains	40 Darling Downs	45 Sydney
26 Northern Plains	41 Darling Downs	46 Sydney
27 Northern Plains	42 Darling Downs	47 Sydney
28 Northern Plains	43 Darling Downs	48 Sydney
29 Northern Plains	44 Darling Downs	49 Sydney
30 Northern Plains	45 Darling Downs	50 Sydney
31 Northern Plains	46 Darling Downs	51 Sydney
32 Northern Plains	47 Darling Downs	52 Sydney
33 Northern Plains	48 Darling Downs	53 Sydney
34 Northern Plains	49 Darling Downs	54 Sydney
35 Northern Plains	50 Darling Downs	55 Sydney
36 Northern Plains	51 Darling Downs	56 Sydney
37 Northern Plains	52 Darling Downs	57 Sydney
38 Northern Plains	53 Darling Downs	58 Sydney
39 Northern Plains	54 Darling Downs	59 Sydney
40 Northern Plains	55 Darling Downs	60 Sydney
41 Northern Plains	56 Darling Downs	61 Sydney
42 Northern Plains	57 Darling Downs	62 Sydney
43 Northern Plains	58 Darling Downs	63 Sydney
44 Northern Plains	59 Darling Downs	64 Sydney
45 Northern Plains	60 Darling Downs	65 Sydney
46 Northern Plains	61 Darling Downs	66 Sydney
47 Northern Plains	62 Darling Downs	67 Sydney
48 Northern Plains	63 Darling Downs	68 Sydney
49 Northern Plains	64 Darling Downs	69 Sydney
50 Northern Plains	65 Darling Downs	70 Sydney
51 Northern Plains	66 Darling Downs	71 Sydney
52 Northern Plains	67 Darling Downs	72 Sydney
53 Northern Plains	68 Darling Downs	73 Sydney
54 Northern Plains	69 Darling Downs	74 Sydney
55 Northern Plains	70 Darling Downs	75 Sydney
56 Northern Plains	71 Darling Downs	76 Sydney
57 Northern Plains	72 Darling Downs	77 Sydney
58 Northern Plains	73 Darling Downs	78 Sydney
59 Northern Plains	74 Darling Downs	79 Sydney
60 Northern Plains	75 Darling Downs	80 Sydney
61 Northern Plains	76 Darling Downs	81 Sydney
62 Northern Plains	77 Darling Downs	82 Sydney
63 Northern Plains	78 Darling Downs	83 Sydney
64 Northern Plains	79 Darling Downs	84 Sydney
65 Northern Plains	80 Darling Downs	85 Sydney
66 Northern Plains	81 Darling Downs	86 Sydney
67 Northern Plains	82 Darling Downs	87 Sydney
68 Northern Plains	83 Darling Downs	88 Sydney
69 Northern Plains	84 Darling Downs	89 Sydney
70 Northern Plains	85 Darling Downs	90 Sydney
71 Northern Plains	86 Darling Downs	91 Sydney
72 Northern Plains	87 Darling Downs	92 Sydney
73 Northern Plains	88 Darling Downs	93 Sydney
74 Northern Plains	89 Darling Downs	94 Sydney
75 Northern Plains	90 Darling Downs	95 Sydney
76 Northern Plains	91 Darling Downs	96 Sydney
77 Northern Plains	92 Darling Downs	97 Sydney
78 Northern Plains	93 Darling Downs	98 Sydney
79 Northern Plains	94 Darling Downs	99 Sydney
80 Northern Plains	95 Darling Downs	100 Sydney
81 Northern Plains	96 Darling Downs	101 Sydney
82 Northern Plains	97 Darling Downs	102 Sydney
83 Northern Plains	98 Darling Downs	103 Sydney
84 Northern Plains	99 Darling Downs	104 Sydney
85 Northern Plains	100 Darling Downs	105 Sydney
86 Northern Plains	101 Darling Downs	106 Sydney
87 Northern Plains	102 Darling Downs	107 Sydney
88 Northern Plains	103 Darling Downs	108 Sydney
89 Northern Plains	104 Darling Downs	109 Sydney
90 Northern Plains	105 Darling Downs	110 Sydney
91 Northern Plains	106 Darling Downs	111 Sydney
92 Northern Plains	107 Darling Downs	112 Sydney
93 Northern Plains	108 Darling Downs	113 Sydney
94 Northern Plains	109 Darling Downs	114 Sydney
95 Northern Plains	110 Darling Downs	115 Sydney
96 Northern Plains	111 Darling Downs	116 Sydney
97 Northern Plains	112 Darling Downs	117 Sydney
98 Northern Plains	113 Darling Downs	118 Sydney
99 Northern Plains	114 Darling Downs	119 Sydney
100 Northern Plains	115 Darling Downs	120 Sydney


COMMONWEALTH OF AUSTRALIA
 METEOROLOGICAL BRANCH · DEPARTMENT OF
 THE INTERIOR · MELBOURNE

MAP OF AUSTRALIA

SHOWING
**AVERAGE ANNUAL
 EVAPORATION**

SCALE MILES 0 100 200 300


FROM EVAPORATION RECORD TO END OF 1946 AND SATURATION DEFICIT RECORDS USING WAITE FORMULA
 $e = 263 S.d.$
 (SEE LETTERPRESS ACCOMPANYING)





COMMONWEALTH OF AUSTRALIA.
BUREAU OF METEOROLOGY MELBOURNE.
MONTHLY DISTRIBUTION OF RAINFALL
OVER AUSTRALIA.

REVISED TO END OF 1939.

ISSUED UNDER THE AUTHORITY OF THE MINISTER OF STATE FOR THE AIR BY
H. N. WARREN
DIRECTOR.



EXPLANATION:- THE GRAPHS SHOW THE AVERAGE RAINFALL IN EACH MONTH OF THE YEAR IN THE VARIOUS RAINFALL DISTRICTS. THE BOUNDARIES OF WHICH ARE SHOWN BY THE BROKEN LINES ON THE MAP. THE NUMBERS ON THE MAP REFER TO THE DISTRICTS. THE CORRESPONDING DISTRICT NAMES ARE GIVEN IN THE INSERT ON PAGE 21.
THE COLUMNS FOR THE SUCCESSIVE MONTHS, JANUARY TO DECEMBER, RUN FROM LEFT TO RIGHT.
THE LENGTH OF A VERTICAL COLUMN REPRESENTS THE AVERAGE MONTHLY RAINFALL ACCORDING TO THE SCALE GIVEN BELOW.




STATE LIBRARY OF VICTORIA

ISSUED UNDER THE AUTHORITY OF THE MINISTER OF STATE FOR THE DEPARTMENT OF AIR
BY H.N. WARREN DIRECTOR OF METEOROLOGICAL SERVICES

9 A.M. LOCAL MEAN TIME WIND ROSES


SHOWING MONTHLY FREQUENCY
DISTRIBUTION OF WIND DIRECTIONS
TO EIGHT POINTS OF THE COMPASS



ISSUED UNDER THE AUTHORITY OF THE MINISTER OF STATE FOR THE DEPARTMENT OF AIR
BY H.N. WARREN DIRECTOR OF METEOROLOGICAL SERVICES

3 P.M. LOCAL MEAN TIME WIND ROSES

SHOWING MONTHLY FREQUENCY
DISTRIBUTION OF WIND DIRECTIONS
TO EIGHT POINTS OF THE COMPASS



The area affected and the period of duration of the longest heat waves in Australia are shown in the map and diagram on page 34.

(v) *Tabulated Data for Selected Climatological Stations in Australia.* The following tables show normal mean temperature, extreme temperature and normal rainfall for each month for selected climatological stations in each State:—

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS: NEW SOUTH WALES.

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
GRAFTON.													
Normal Mean Temperature—													
Maximum °F.	89.1	87.9	85.2	81.7	76.1	70.9	70.6	73.3	78.6	82.6	85.7	88.0	80.8
Minimum ..	66.4	66.3	63.7	57.9	51.0	45.7	43.3	45.1	50.4	56.3	61.3	64.5	56.0
Extreme Temperature—													
Maximum .. °F.	114.0	113.0	108.0	97.0	91.0	88.0	87.5	95.0	99.0	105.0	111.0	113.5	114.0
Minimum ..	50.0	50.0	41.0	33.0	33.0	28.0	24.9	24.0	32.0	35.0	43.0	45.0	24.0
Normal Rainfall ins.	4.56	4.19	3.72	3.15	2.77	2.44	2.03	0.93	1.83	2.23	3.31	3.52	34.68

ARMIDALE.													
Normal Mean Temperature—													
Maximum °F.	80.8	79.5	75.3	68.4	61.2	55.3	54.0	57.2	63.8	70.4	76.1	79.3	68.4
Minimum ..	56.5	55.8	52.1	45.6	39.2	34.9	33.8	34.4	38.9	45.1	50.3	54.3	45.1
Extreme Temperature—													
Maximum .. °F.	103.4	95.0	94.0	86.2	80.0	76.0	68.2	78.2	83.0	90.5	97.5	99.8	103.4
Minimum ..	40.0	38.0	31.0	25.0	20.0	17.0	14.0	18.0	22.0	26.0	32.0	36.0	14.0
Normal Rainfall ins.	3.88	2.81	2.26	1.87	1.46	2.33	2.11	1.54	2.09	2.35	2.87	3.41	28.98

PORT MACQUARIE.													
Normal Mean Temperature—													
Maximum °F.	78.6	78.7	77.1	73.2	68.8	64.9	64.0	65.8	68.5	71.2	74.0	76.4	71.8
Minimum ..	64.4	64.3	61.8	56.8	50.8	46.3	44.8	45.4	49.2	54.8	59.0	62.5	55.0
Extreme Temperature—													
Maximum .. °F.	104.0	105.8	97.0	92.8	84.0	79.6	84.0	91.3	89.4	97.4	104.0	98.8	105.8
Minimum ..	51.0	48.0	43.5	40.0	34.6	30.5	29.5	31.0	32.5	38.0	41.2	48.0	29.5
Normal Rainfall ins.	4.89	6.48	6.45	7.37	5.76	5.08	4.35	2.64	3.55	3.67	3.22	3.98	57.44

BROKEN HILL.													
Normal Mean Temperature—													
Maximum °F.	90.5	90.2	84.6	74.7	66.7	59.9	59.5	63.6	70.1	77.3	83.3	88.6	75.8
Minimum ..	64.5	64.7	60.1	52.6	47.1	42.5	41.2	43.2	47.6	52.9	58.1	62.6	53.1
Extreme Temperature—													
Maximum .. °F.	114.9	115.9	113.9	99.9	87.8	79.0	80.0	84.0	94.0	103.9	110.9	113.9	115.9
Minimum ..	45.0	42.0	40.0	34.0	30.5	27.0	28.5	29.0	33.0	36.0	40.9	41.8	27.0
Normal Rainfall ins.	0.59	0.96	0.56	0.65	0.94	0.92	0.72	0.63	0.67	0.84	0.86	0.86	9.20

DUBBO.													
Normal Mean Temperature—													
Maximum °F.	92.1	91.3	85.7	76.9	68.0	61.0	59.7	63.5	70.3	78.5	85.3	89.6	76.8
Minimum ..	63.8	63.8	58.9	50.8	43.5	39.3	37.5	38.3	42.7	49.3	56.4	61.3	50.5
Extreme Temperature—													
Maximum .. °F.	114.9	113.9	104.5	97.9	90.9	79.9	77.8	87.0	92.9	104.9	110.0	115.4	115.4
Minimum ..	40.9	35.7	37.7	30.0	23.4	19.9	16.9	17.9	20.9	27.9	30.9	37.9	16.9
Normal Rainfall ins.	2.00	1.49	1.99	1.77	1.44	2.24	1.87	1.51	1.31	1.49	1.93	1.87	20.91

CHAPTER II.—PHYSIOGRAPHY.

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS:
NEW SOUTH WALES—*continued.*

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
NEWCASTLE.													
Normal Mean Temperature—													
Maximum °F.	77.6	77.7	76.1	72.2	67.2	62.9	61.7	64.0	68.1	71.3	73.9	76.0	70.7
Minimum ..	66.6	67.1	64.7	59.5	53.7	49.5	47.7	48.8	52.6	57.2	61.3	64.3	57.7
Extreme Temperature—													
Maximum .. °F.	112.0	105.3	101.5	94.9	85.0	80.0	79.4	88.3	96.4	100.0	105.0	108.0	112.0
Minimum ..	54.5	54.0	50.0	42.0	41.0	38.0	37.5	37.0	39.0	42.0	47.8	49.0	37.0
Normal Rainfall ins.	3.01	2.96	3.84	5.33	4.64	3.66	4.44	2.34	2.97	2.51	2.21	3.45	41.36
BATHURST.													
Normal Mean Temperature—													
Maximum °F.	83.9	83.7	78.8	69.9	62.1	55.2	53.9	57.4	64.2	70.9	76.7	81.6	69.9
Minimum ..	55.6	55.5	51.0	44.0	38.2	34.8	34.0	34.5	38.1	43.4	48.4	53.5	44.2
Extreme Temperature—													
Maximum .. °F.	112.9	106.4	100.2	90.0	80.0	71.0	70.0	76.4	86.0	96.0	103.5	107.7	112.9
Minimum ..	37.0	35.0	30.0	22.0	20.0	15.7	13.0	18.7	21.0	25.0	31.0	35.0	13.0
Normal Rainfall ins.	2.18	1.73	1.99	1.51	1.39	1.86	2.07	1.68	1.52	2.10	2.04	2.49	22.56
LEETON.													
Normal Mean Temperature—													
Maximum °F.	88.9	88.7	82.6	72.6	64.8	57.7	56.8	60.3	66.8	73.8	81.2	86.4	73.4
Minimum ..	63.2	63.4	59.0	51.2	45.0	40.5	38.9	40.5	44.1	49.7	55.6	60.8	51.0
Extreme Temperature—													
Maximum .. °F.	117.0	110.5	107.0	94.5	82.4	74.5	72.0	81.9	92.5	103.5	107.0	112.0	117.0
Minimum ..	44.0	41.2	40.0	33.0	29.9	24.9	25.3	25.0	26.5	34.0	35.5	41.8	24.9
Normal Rainfall ins.	1.22	0.86	1.03	1.47	1.38	1.84	1.36	1.67	1.31	1.49	1.26	1.24	16.13
JERVIS BAY.(a)													
Normal Mean Temperature—													
Maximum °F.	74.5	75.0	73.3	68.8	64.4	60.2	58.9	61.0	64.3	67.7	70.2	73.0	67.6
Minimum ..	63.0	64.0	62.7	58.5	53.8	50.5	48.6	49.5	52.2	55.3	58.3	61.3	56.5
Extreme Temperature—													
Maximum .. °F.	109.0	100.0	98.0	89.0	83.0	77.0	75.0	80.0	87.0	96.0	96.0	100.0	109.0
Minimum ..	43.0	50.0	46.0	42.0	39.0	37.0	33.0	31.0	40.0	41.0	47.0	42.0	31.0
Normal Rainfall ins.	4.11	3.25	4.46	5.07	5.22	4.32	4.89	2.72	3.06	2.57	2.64	3.82	46.13
ALBURY.													
Normal Mean Temperature—													
Maximum °F.	89.9	90.4	84.3	73.6	64.9	57.4	56.4	60.4	67.2	73.8	81.3	87.4	73.9
Minimum ..	59.8	60.2	55.2	47.8	42.3	39.3	38.2	39.9	43.2	47.7	52.9	57.5	48.7
Extreme Temperature—													
Maximum .. °F.	117.3	114.3	107.3	94.8	83.0	76.0	74.0	79.0	94.8	101.5	107.0	112.6	117.3
Minimum ..	39.0	42.0	39.0	30.0	28.0	25.7	25.0	26.0	29.0	30.0	33.0	40.0	25.0
Normal Rainfall ins.	1.59	1.95	1.81	1.96	2.28	3.32	2.91	3.01	2.30	2.52	1.76	2.25	27.66
COOMA.													
Normal Mean Temperature—													
Maximum °F.	78.8	79.0	73.8	65.0	57.3	50.9	50.4	54.5	61.0	67.5	72.6	77.2	65.7
Minimum ..	52.2	52.4	48.3	41.7	35.3	31.7	30.2	31.5	36.1	40.9	45.8	50.1	41.3
Extreme Temperature—													
Maximum .. °F.	112.0	107.0	104.6	92.7	77.7	69.2	72.9	75.7	86.9	95.7	102.1	110.0	112.0
Minimum ..	29.8	33.0	28.2	22.8	13.0	13.4	11.0	12.0	14.3	22.0	25.8	28.8	11.0
Normal Rainfall ins.	2.34	1.86	1.88	1.46	1.11	1.22	1.27	0.98	1.27	1.51	1.72	2.23	18.85

(a) Australian Capital Territory.

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS: VICTORIA.

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
MILDURA.													
Normal Mean Temperature—													
Maximum °F.	89.8	90.0	84.4	74.5	66.9	60.4	59.5	63.9	69.9	76.5	83.2	88.2	75.6
Minimum ..	61.0	61.7	57.2	50.5	45.6	41.3	40.5	42.5	46.1	50.9	55.4	59.6	51.0
Extreme Temperature—													
Maximum .. °F.	123.5	118.0	112.0	99.0	90.0	80.0	78.0	86.9	95.0	104.0	113.0	121.5	123.5
Minimum ..	40.0	43.0	37.0	34.0	27.0	26.0	24.0	29.0	29.0	34.0	35.0	40.0	24.0
Normal Rainfall ins.	0.73	0.90	0.70	0.55	1.01	1.05	0.91	1.01	0.96	1.00	0.84	0.71	10.37
BENALLA.													
Normal Mean Temperature—													
Maximum °F.	87.6	88.7	82.4	72.3	64.3	56.5	55.7	58.9	65.1	72.4	79.5	84.8	72.3
Minimum ..	58.9	59.6	55.1	48.0	42.6	39.1	38.2	39.7	43.6	48.3	52.1	56.5	48.5
Extreme Temperature—													
Maximum .. °F.	114.0	112.0	108.0	97.0	89.0	70.0	70.0	76.0	88.0	102.0	104.0	111.0	114.0
Minimum ..	40.0	37.0	36.0	32.0	26.0	25.0	27.0	27.9	30.0	33.0	36.0	38.0	25.0
Normal Rainfall ins.	1.58	1.66	1.60	1.99	2.30	3.09	2.78	2.72	2.36	2.41	1.65	1.77	25.91
BENDIGO.													
Normal Mean Temperature—													
Maximum °F.	83.0	83.9	78.1	68.4	61.3	54.8	54.2	57.0	62.5	68.9	75.2	80.5	69.0
Minimum ..	56.5	58.3	54.0	48.2	43.7	40.7	39.4	40.2	43.0	46.7	50.9	54.9	48.0
Extreme Temperature—													
Maximum .. °F.	117.4	111.6	104.7	94.7	80.0	77.3	73.0	75.7	90.0	99.7	106.5	111.5	117.4
Minimum ..	37.0	40.0	38.0	33.4	27.3	23.9	23.5	26.0	29.0	32.0	35.0	37.0	23.5
Normal Rainfall ins.	1.14	1.50	1.27	1.49	1.97	2.26	2.21	2.11	2.04	1.70	1.25	1.33	20.27
HORSHAM.													
Normal Mean Temperature—													
Maximum °F.	85.1	86.3	80.2	70.7	63.0	56.6	56.0	59.0	64.1	70.2	77.2	82.7	70.9
Minimum ..	55.2	55.9	51.9	47.0	42.9	40.2	38.8	39.9	41.9	45.1	49.6	53.2	46.8
Extreme Temperature—													
Maximum .. °F.	120.0	113.0	108.0	97.0	87.0	74.0	71.0	78.0	94.0	100.0	108.0	115.0	120.0
Minimum ..	39.0	37.0	35.0	31.0	25.0	22.0	21.0	24.0	24.0	25.0	29.0	34.0	21.0
Normal Rainfall ins.	0.75	1.21	0.74	1.23	1.78	1.98	1.89	1.90	1.98	1.48	1.26	1.37	17.57
BALLARAT.													
Normal Mean Temperature—													
Maximum °F.	75.7	76.9	71.6	63.0	56.3	50.4	49.8	52.5	57.1	62.4	67.4	72.5	63.0
Minimum ..	50.5	52.9	50.1	45.8	42.6	39.5	38.4	39.4	41.2	43.6	46.0	49.3	44.9
Extreme Temperature—													
Maximum .. °F.	108.5	104.9	102.1	91.2	75.0	63.0	63.0	69.6	83.0	92.5	100.0	102.0	108.5
Minimum ..	36.0	36.2	31.0	31.0	27.0	23.0	26.0	26.3	26.6	29.0	31.5	35.0	23.0
Normal Rainfall ins.	1.26	1.79	1.83	2.13	2.43	2.67	2.68	2.92	2.84	2.41	2.08	2.34	27.38
BAIRNSDALE.													
Normal Mean Temperature—													
Maximum °F.	75.3	76.1	73.0	67.5	62.5	57.5	57.0	59.5	63.2	67.5	70.6	74.0	67.0
Minimum ..	53.5	54.5	51.7	46.9	42.5	38.8	38.1	39.6	42.7	46.1	49.0	52.4	46.3
Extreme Temperature—													
Maximum .. °F.	112.0	109.0	105.5	95.0	86.0	75.0	76.0	84.0	92.6	101.0	103.0	111.0	112.0
Minimum ..	35.0	39.0	32.0	29.0	25.0	22.0	21.0	19.0	26.0	27.0	30.0	32.0	19.0
Normal Rainfall ins.	2.48	2.09	2.64	2.02	1.59	2.16	2.06	1.73	2.08	2.68	2.19	2.63	26.35

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS : QUEENSLAND.

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
CAIRNS.													
Normal Mean Temperature—													
Maximum °F.	89.7	89.0	87.1	84.9	81.6	78.8	78.1	79.5	82.6	85.6	87.9	89.7	84.5
Minimum ..	74.2	73.9	72.6	70.0	66.2	63.5	61.0	61.1	63.8	67.4	70.4	72.9	68.1
Extreme Temperature—													
Maximum .. °F.	109.8	108.0	100.0	94.5	92.0	92.1	95.1	98.1	94.1	98.1	99.1	105.0	109.8
Minimum ..	63.5	64.0	59.8	57.0	52.2	44.6	43.0	43.2	46.0	54.5	52.0	60.2	43.0
Normal Rainfall ins.	16.51	17.00	17.59	10.76	4.37	2.87	1.56	1.46	1.43	2.40	3.05	7.35	86.35
TOWNSVILLE.													
Normal Mean Temperature—													
Maximum °F.	87.3	87.0	86.6	84.7	81.2	77.3	76.0	77.6	80.3	83.1	85.2	87.0	82.8
Minimum ..	76.2	75.6	73.9	70.6	65.4	61.9	59.8	61.5	65.8	70.5	73.8	75.6	69.2
Extreme Temperature—													
Maximum .. °F.	102.4	110.2	98.0	97.0	88.5	86.5	85.0	89.0	92.5	94.7	99.5	101.1	110.2
Minimum ..	68.2	64.5	65.4	53.9	49.7	47.0	45.3	48.0	52.0	60.2	64.2	66.0	45.3
Normal Rainfall ins.	10.03	9.90	5.15	2.57	0.92	1.41	0.77	0.60	0.49	1.19	2.03	4.63	39.69
CLONGURRY.													
Normal Mean Temperature—													
Maximum °F.	98.7	96.3	94.6	89.9	82.9	77.3	76.4	81.4	88.4	95.1	98.6	100.4	90.0
Minimum ..	76.5	75.4	73.0	66.9	59.7	54.1	51.5	54.3	61.0	68.2	73.5	76.2	65.9
Extreme Temperature—													
Maximum .. °F.	127.5	115.5	110.5	108.0	98.5	99.1	96.0	102.5	106.0	112.0	118.5	125.5	127.5
Minimum ..	59.3	58.0	53.3	48.0	41.3	32.0	34.2	34.5	40.5	49.8	54.0	50.0	32.0
Normal Rainfall ins.	4.73	3.96	1.86	0.62	0.48	0.80	0.23	0.12	0.15	0.44	1.59	1.90	16.88
MACKAY.													
Normal Mean Temperature—													
Maximum °F.	86.2	85.5	83.6	80.7	76.1	72.2	71.0	72.8	77.0	81.3	83.9	86.2	79.7
Minimum ..	73.6	73.2	71.2	66.6	60.8	56.2	53.4	54.8	59.9	65.5	69.4	72.3	64.7
Extreme Temperature—													
Maximum .. °F.	99.8	99.4	98.0	94.0	88.8	85.9	86.0	87.0	92.0	97.0	97.5	99.9	99.9
Minimum ..	60.1	60.3	56.0	49.2	41.6	37.0	35.1	36.1	39.6	44.0	46.6	60.0	35.1
Normal Rainfall ins.	13.56	12.65	10.95	4.64	3.36	2.75	1.57	1.12	1.14	1.55	3.12	6.75	63.16
LONGREACH.													
Normal Mean Temperature—													
Maximum °F.	99.6	96.9	94.1	87.8	80.4	74.3	73.2	77.9	85.4	92.8	97.0	99.7	88.3
Minimum ..	73.3	71.7	68.1	60.1	52.1	46.7	44.3	46.5	53.7	61.5	67.5	71.5	59.8
Extreme Temperature—													
Maximum .. °F.	118.3	113.4	113.0	103.0	96.8	92.0	92.0	96.8	104.2	109.2	114.5	115.6	118.3
Minimum ..	43.9	55.1	48.2	38.1	35.1	26.7	26.0	31.0	31.0	39.0	41.0	43.9	26.7
Normal Rainfall ins.	2.31	3.12	2.10	1.01	0.52	0.94	0.80	0.30	0.52	0.84	1.26	1.82	15.34
ROCKHAMPTON.													
Normal Mean Temperature—													
Maximum °F.	90.0	88.7	87.2	84.2	79.3	74.4	73.7	76.7	81.7	85.9	88.5	90.0	83.4
Minimum ..	72.3	72.1	69.8	64.8	58.3	54.0	51.2	52.9	58.3	63.8	68.0	70.9	63.0
Extreme Temperature—													
Maximum .. °F.	106.8	105.2	104.8	98.0	94.3	88.2	88.8	95.9	100.2	102.8	107.9	111.6	111.6
Minimum ..	60.0	60.8	59.3	43.4	39.8	32.7	34.6	30.3	39.8	43.0	54.0	59.2	32.7
Normal Rainfall ins.	6.70	7.28	3.54	2.66	1.26	2.80	1.77	0.82	0.94	1.99	2.63	4.97	37.36

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS:
QUEENSLAND—*continued.*

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.	
CHARLEVILLE.														
Normal Mean Temperature—														
Maximum °F.	97.6	96.1	91.7	84.5	76.4	69.3	68.3	72.9	80.4	88.2	93.6	96.4	84.6	
Minimum .. "	70.8	70.1	65.1	55.7	47.2	42.3	40.1	42.1	49.0	57.7	64.4	68.5	56.1	
Extreme Temperature—														
Maximum .. °F.	116.6	115.0	110.0	101.8	92.0	87.8	86.5	93.7	102.0	109.8	117.0	118.0	118.0	
Minimum .. "	52.4	50.0	41.0	34.0	27.5	23.0	23.0	24.0	29.0	34.5	40.0	48.0	23.0	
Normal Rainfall	ins.	2.65	2.30	1.54	0.95	0.69	1.46	1.32	0.75	0.95	1.02	1.68	2.60	17.97
TOOWOOMBA.														
Normal Mean Temperature—														
Maximum °F.	82.7	81.0	78.2	73.5	67.2	62.0	61.1	64.5	70.5	76.2	80.2	82.2	73.3	
Minimum .. "	61.2	61.0	58.7	52.5	46.6	42.4	40.7	41.9	46.8	52.3	56.6	59.5	51.7	
Extreme Temperature—														
Maximum .. °F.	103.6	100.7	99.0	88.2	84.2	80.5	78.2	86.0	89.5	96.0	101.0	105.0	105.0	
Minimum .. "	45.5	46.0	33.0	31.0	29.6	22.0	22.5	25.0	30.0	32.4	39.4	42.8	22.0	
Normal Rainfall	ins.	5.15	4.29	3.36	2.62	1.85	2.54	2.06	1.16	1.69	2.39	3.34	4.74	35.19

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS: SOUTH AUSTRALIA.

STREAKY BAY.

Normal Mean Temperature—														
Maximum °F.	84.8	85.2	81.8	74.2	67.9	62.0	61.0	63.3	67.7	73.2	78.6	82.0	73.5	
Minimum .. "	60.1	60.5	58.5	54.1	51.4	48.2	46.9	47.8	49.2	52.4	55.6	58.3	53.6	
Extreme Temperature—														
Maximum .. °F.	114.2	114.2	109.0	96.0	88.3	79.0	73.0	83.0	91.0	104.2	113.8	117.0	117.0	
Minimum .. "	46.2	44.8	43.5	41.0	34.0	31.0	31.2	32.2	33.9	38.0	39.5	42.5	31.0	
Normal Rainfall	ins.	0.30	0.68	0.57	0.76	1.72	2.64	2.29	2.19	1.31	1.01	0.69	0.56	14.62

PORT PIRIE.

Normal Mean Temperature—														
Maximum °F.	89.2	89.6	86.4	76.8	69.5	62.8	61.7	64.8	71.2	77.1	82.9	86.2	76.5	
Minimum .. "	62.9	63.2	61.1	55.2	50.9	46.5	45.5	46.5	49.2	53.4	57.4	60.9	54.4	
Extreme Temperature—														
Maximum .. °F.	117.1	113.0	111.0	99.0	85.0	77.0	76.0	83.0	95.0	103.0	109.0	114.2	117.1	
Minimum .. "	48.2	48.2	47.0	41.0	36.0	30.0	31.0	33.0	35.0	37.0	43.0	46.0	30.0	
Normal Rainfall	ins.	0.75	0.83	0.70	0.78	1.40	1.54	1.25	1.48	1.23	1.17	0.92	0.94	12.99

YONGALA.

Normal Mean Temperature—														
Maximum °F.	85.9	85.8	80.7	70.5	62.4	55.5	54.5	57.6	63.8	71.2	78.3	83.5	70.8	
Minimum .. "	55.8	56.3	51.7	45.1	40.9	37.6	36.1	36.9	39.4	43.4	49.0	53.7	45.5	
Extreme Temperature—														
Maximum .. °F.	111.2	107.6	105.0	95.0	83.0	71.8	72.4	79.2	91.0	98.2	104.0	107.0	111.2	
Minimum .. "	38.0	39.0	34.0	28.4	23.5	19.0	19.0	24.0	25.8	24.0	30.2	35.0	19.0	
Normal Rainfall	ins.	0.80	0.86	0.60	0.88	1.37	1.53	1.62	1.87	1.54	1.23	1.10	1.16	14.56

MT. GAMBIER.

Normal Mean Temperature—														
Maximum °F.	74.2	75.9	72.7	66.5	61.4	57.0	56.2	58.1	61.1	65.0	68.3	71.9	65.7	
Minimum .. "	53.5	54.8	52.4	49.5	46.4	43.5	42.4	43.1	45.1	46.9	49.6	52.0	48.3	
Extreme Temperature—														
Maximum .. °F.	112.6	109.6	106.3	98.2	82.6	72.0	70.5	77.0	89.0	95.0	104.0	107.6	112.6	
Minimum .. "	33.0	34.0	33.0	29.8	26.2	23.4	23.7	27.5	28.4	30.4	31.0	34.0	23.4	
Normal Rainfall	ins.	0.93	1.22	1.17	2.14	2.90	3.55	3.49	3.44	2.91	2.10	1.56	1.45	26.

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS : WESTERN AUSTRALIA.

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
--------------	------	------	------	------	------	-------	-------	------	-------	------	------	------	-------

WYNDHAM.

Normal Mean Temperature—														
Maximum	°F.	95.9	95.5	95.3	94.7	90.1	85.8	85.0	88.5	93.5	96.9	98.5	97.6	93.1
Minimum	"	80.2	79.7	79.5	77.2	72.4	68.0	66.2	69.5	74.8	79.7	81.4	81.2	75.8
Extreme Temperature—														
Maximum ..	°F.	113.5	108.3	106.0	104.6	102.4	97.5	96.0	102.0	106.0	110.2	111.6	109.5	113.5
Minimum ..	"	67.5	62.0	65.0	63.5	55.2	50.0	48.0	56.0	60.1	65.0	68.0	67.0	48.0
Normal Rainfall	ins.	6.79	6.30	5.22	0.50	0.14	0.20	0.08	0.02	0.05	0.38	1.55	3.92	25.15

GERALDTON.

Normal Mean Temperature—														
Maximum	°F.	84.5	85.2	83.6	80.5	74.2	69.7	67.7	68.8	71.4	73.6	78.5	82.0	76.6
Minimum	"	66.3	66.5	65.0	60.9	56.9	53.8	51.7	52.1	53.0	55.4	60.0	63.4	58.7
Extreme Temperature—														
Maximum ..	°F.	112.0	115.5	110.8	102.8	93.8	83.8	81.0	86.0	96.5	104.6	108.8	113.0	115.5
Minimum ..	"	48.0	51.0	47.0	41.8	38.6	33.6	33.4	37.3	38.3	41.0	44.0	48.0	33.4
Normal Rainfall	ins.	0.30	0.42	0.78	0.89	2.58	4.84	3.77	2.57	1.21	0.79	0.27	0.16	18.58

KALGOORLIE.

Normal Mean Temperature—														
Maximum	°F.	93.2	93.0	86.3	78.4	70.1	63.6	62.5	66.0	73.6	79.0	86.3	91.1	78.6
Minimum	"	64.2	64.4	61.3	55.2	48.9	44.6	42.9	43.9	48.2	52.7	58.3	62.3	53.9
Extreme Temperature—														
Maximum ..	°F.	114.4	115.0	111.0	102.5	92.0	81.8	81.0	87.0	96.0	102.3	110.6	113.0	115.0
Minimum ..	"	47.1	48.0	41.6	37.0	34.6	31.0	30.0	30.0	31.6	33.4	38.2	46.0	30.0
Normal Rainfall	ins.	0.69	0.65	1.26	0.97	0.98	0.98	0.81	0.88	0.38	0.58	0.61	0.67	9.46

COLLIE.

Normal Mean Temperature—														
Maximum	°F.	86.4	85.7	80.4	74.3	65.9	61.3	59.8	61.0	64.8	68.8	77.2	83.0	72.4
Minimum	"	55.6	54.9	52.5	47.1	42.9	40.4	39.1	39.8	42.5	45.3	49.7	53.1	46.9
Extreme Temperature—														
Maximum ..	°F.	109.0	110.2	105.3	98.0	86.8	76.0	73.0	79.0	86.6	96.4	101.8	106.2	110.2
Minimum ..	"	37.7	35.2	32.3	29.6	28.0	24.8	25.0	26.2	28.0	31.0	32.6	35.0	24.8
Normal Rainfall	ins.	0.67	0.70	0.98	1.85	5.24	6.91	7.84	6.10	4.44	3.06	1.12	0.69	39.60

ALBANY.

Normal Mean Temperature—														
Maximum	°F.	73.8	74.2	72.3	70.3	65.9	62.2	60.9	61.7	63.6	65.7	69.2	72.0	67.6
Minimum	"	58.5	58.8	57.5	54.5	50.7	47.8	46.3	46.6	48.3	50.0	53.6	56.5	54.4
Extreme Temperature—														
Maximum ..	°F.	106.0	112.6	105.4	99.6	95.3	76.2	73.5	81.0	87.0	97.2	106.0	106.0	112.6
Minimum ..	"	42.3	41.0	38.7	39.5	35.1	35.0	32.2	34.3	34.0	36.2	40.6	41.2	32.2
Normal Rainfall	ins.	1.36	1.03	1.78	2.93	5.30	5.44	6.00	5.42	4.25	3.28	1.65	1.23	39.67

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS : TASMANIA.

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
BURNIE.													
Normal Mean Temperature—													
Maximum °F.	66.7	66.9	64.6	61.1	57.9	54.3	52.9	53.7	55.7	57.9	61.6	65.3	59.9
Minimum ..	50.9	52.9	49.4	47.4	44.5	43.2	40.5	41.3	41.8	43.7	47.1	49.7	46.1
Extreme Temperature—													
Maximum .. °F.	83.5	80.3	80.0	71.3	70.5	60.4	59.8	61.3	63.0	72.7	80.4	85.0	85.0
Minimum ..	39.5	38.7	36.0	35.5	31.3	32.8	30.5	30.0	31.0	32.0	36.0	36.5	30.0
Normal Rainfall ins.	1.52	1.73	1.88	3.23	3.53	4.55	4.92	4.88	3.83	3.65	2.62	2.65	38.99

LAUNCESTON.													
Normal Mean Temperature—													
Maximum °F.	75.8	76.7	72.0	65.3	59.5	54.6	53.7	56.3	60.0	64.2	69.2	73.1	65.0
Minimum ..	52.1	52.7	49.7	45.3	41.1	38.4	36.9	38.4	41.4	44.1	47.2	50.3	44.8
Extreme Temperature—													
Maximum .. °F.	100.0	101.0	98.5	84.0	74.8	66.2	66.2	68.0	75.0	88.8	92.0	97.6	101.0
Minimum ..	34.0	33.7	31.0	27.0	24.0	22.0	21.0	24.5	24.0	25.0	32.0	31.5	21.0
Normal Rainfall ins.	1.52	1.49	1.71	2.38	2.71	3.11	3.10	3.12	2.81	2.67	1.82	2.12	28.56

ZEEHAN.													
Normal Mean Temperature—													
Maximum °F.	66.3	68.6	65.2	59.9	56.0	52.3	51.6	53.0	55.9	58.9	61.4	64.7	59.5
Minimum ..	48.0	49.4	47.3	45.1	42.4	39.2	38.2	39.7	41.0	42.8	44.6	46.8	43.7
Extreme Temperature—													
Maximum .. °F.	94.7	99.2	92.5	81.9	74.6	67.3	62.6	69.3	80.0	85.9	94.0	97.0	99.2
Minimum ..	32.2	30.7	28.0	25.0	23.0	20.5	21.1	22.6	23.0	26.6	30.0	31.2	20.5
Normal Rainfall ins.	5.75	4.37	5.77	8.25	8.69	9.21	9.90	10.30	9.34	8.59	7.43	6.46	94.06

SWANSEA.													
Normal Mean Temperature—													
Maximum °F.	70.9	71.4	68.7	63.9	59.4	55.3	54.7	56.5	59.9	63.3	66.8	69.2	63.3
Minimum ..	52.1	53.0	50.3	46.6	42.4	40.0	38.9	39.5	42.3	44.9	47.9	50.6	45.7
Extreme Temperature—													
Maximum .. °F.	104.1	100.7	99.7	85.0	83.9	67.0	67.0	72.6	80.0	92.0	95.0	100.2	104.1
Minimum ..	35.4	38.0	27.0	31.4	27.0	24.0	26.4	25.0	27.2	29.0	32.0	35.0	24.0
Normal Rainfall ins.	1.60	1.89	2.48	2.30	1.65	2.52	1.97	1.37	1.51	2.17	1.84	2.49	23.79

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS : NORTHERN TERRITORY.

DARWIN.

Normal Mean Temperature—													
Maximum °F.	89.9	89.8	90.2	91.9	90.1	87.5	86.6	88.5	91.0	92.6	93.2	92.0	90.3
Minimum ..	77.3	77.1	77.1	75.9	72.6	69.5	67.8	69.7	73.9	77.2	78.2	78.1	74.5
Extreme Temperature—													
Maximum .. °F.	100.0	100.9	102.0	104.0	102.3	98.6	98.0	98.0	102.0	104.9	103.3	102.0	104.9
Minimum ..	68.0	63.0	68.0	65.7	59.2	55.3	55.8	58.1	63.0	68.7	66.8	69.4	55.3
Normal Rainfall ins.	16.18	12.37	11.18	3.08	0.33	0.09	0.01	0.02	0.60	1.93	4.32	8.57	58.68

TABULATED DATA FOR SELECTED CLIMATOLOGICAL STATIONS :
NORTHERN TERRITORY—*continued.*

Particulars.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Year.
DALY WATERS.													
Normal Mean Temperature—													
Maximum .. °F.	97.5	96.6	94.9	93.2	88.0	84.0	83.9	89.1	95.6	100.7	102.1	101.1	93.9
Minimum .. °F.	75.6	74.4	72.6	67.1	61.0	56.9	54.0	56.4	63.9	71.0	74.8	75.9	67.0
Extreme Temperature—													
Maximum .. °F.	113.0	111.0	110.3	106.2	101.0	97.8	98.0	102.4	108.1	112.0	113.0	116.1	116.1
Minimum .. °F.	61.2	55.0	55.2	49.0	43.0	35.1	30.2	39.0	41.2	48.3	56.5	61.2	30.2
Normal Rainfall .. ins.	6.77	5.11	4.24	0.94	0.27	0.14	0.03	0.03	0.12	0.68	2.41	3.98	24.74
ALICE SPRINGS.													
Normal Mean Temperature—													
Maximum .. °F.	95.3	94.7	90.1	81.3	73.3	67.1	66.9	72.5	80.1	87.6	91.7	94.7	82.9
Minimum .. °F.	69.8	68.3	62.8	53.8	46.0	41.2	38.9	43.2	49.2	58.5	64.0	68.2	55.3
Extreme Temperature—													
Maximum .. °F.	116.0	114.0	113.0	102.8	101.0	87.0	88.0	96.4	99.7	113.1	115.0	117.0	117.0
Minimum .. °F.	50.0	48.0	39.0	35.4	27.0	22.0	19.0	25.0	30.0	36.4	40.0	46.0	19.0
Normal Rainfall .. ins.	1.74	1.32	1.09	0.39	0.60	0.52	0.29	0.31	0.28	0.71	1.15	1.53	9.93

(vi) *Frosts.** The Observer's Handbook of the Meteorological Office, London, gives the following definition:—"Injury to the tissues of growing plants is not caused until the temperature has fallen considerably below the freezing point of water (32° F.) and a 'ground frost' is regarded as having occurred when the thermometer on the grass has fallen to 30.4° F. or below".

In Australia this definition is adopted for stations equipped with terrestrial minimum thermometers. However, these are few in number, so although many rainfall observers record "hoar frost" when seen, for statistical purposes a screen temperature of 36° F. is taken as indicating light frosts at ground level. For heavy frosts a screen temperature of 32° F. is taken.

In America a "killing" frost is defined as a frost "that is generally destructive of vegetation". A "black frost" is the phenomenon arising out of a combination of low temperature and low humidity causing rupturing of plant cells by expansion, when freezing takes place, of the water which they contain, though frost crystals are not formed on the ground.

The parts of Australia most subject to low temperature are the eastern highlands from about Omeo in Victoria northward to Cambooya and Bybera in Queensland. Most stations in this region experience more than ten nights per month with readings of 32° F. or under for three to five months of the year. In Tasmania, districts on the Central Plateau are subject to such conditions for three to six months of the year. Minimum temperatures of 32° F. are comparatively infrequent in Western Australia except in parts of the south and south-west. In South Australia the Yongala district is much more subject to such temperatures than other parts of the State. Much of the south-east of Queensland has a higher frequency of such readings than South Australia. Generally speaking, the frequency is controlled mainly by altitude, latitude and, to a lesser degree, by proximity to the sea.

Frosts may occur within a few miles of the coastline over the whole continent, except in the Northern Territory and a considerable area of Northern Queensland. Regions subject to frost in all months of the year comprise portions of the tablelands of New South Wales, the Eastern Highlands and parts of the Central Divide and Western district in Victoria, practically the whole of Tasmania and a small area in the south-west of Western Australia.

* Foley, J. C. Frost in the Australian Region (Bull. 32, 1945).

A map showing the average annual number of frost-free days (i.e. days on which the temperature does not fall below 36° F.) appears on page 35.

Over most of the interior of the continent and on the Highlands in Queensland as far north as the Atherton Plateau frosts appear in April and end in September, but they are infrequent in these months. Minimum temperatures of 32° F. are experienced in most of the sub-tropical interior in June and July.

3. **Humidity.**—After temperature, humidity is the most important element of climate, particularly as regards its effects on human comfort, rainfall supply, and conservation and related problems.

In this publication the humidity of the air has been expressed by the relative humidity, which is the quotient of the vapour pressure divided by the saturation vapour pressure and multiplied by one hundred. The mean 9 a.m. relative humidity, as well as its highest and lowest recorded mean values at 9 a.m., are shown in the tables of climatological data for the capital cities (par. 13). The mean monthly vapour pressure has also been added to these tables.

The annual curve of vapour pressure derived from the normal monthly values for this element is comparable with the maximum and minimum temperature curves, but the relative humidities consisting as they do of the extremes for each month do not show the normal annual fluctuation which would be approximately midway between the extremes.

The order of stations in descending values of 9 a.m. vapour pressure is Darwin, Brisbane, Sydney, Perth, Melbourne, Adelaide, Canberra, Hobart and Alice Springs, while the relative humidity diminishes in the order,° Sydney, Canberra, Melbourne, Darwin, Hobart, Brisbane, Perth, Adelaide and Alice Springs.

Further reference to humidity will be found in the section on effective temperature (page 30).

4. **Evaporation.**—(i) *General.* The rate and quantity of evaporation in any territory is influenced by the prevailing temperature, and by atmospheric humidity, pressure and wind movement. In Australia the question is, perhaps, of more than ordinary importance, since in its drier regions water has often to be conserved in "tanks" and dams. The magnitude of the economic loss by evaporation will be appreciated from the map reproduced herein (see page 36) which shows that the yearly amount varies from about 20 inches over Western Tasmania to more than 100 inches over the central and north-western parts of Australia. Over an area of 70 per cent. of the continent, comprising most inland districts and extending to the coast in the North-West and Eucla divisions of Western Australia, during no month of the year does the rainfall exceed the evaporation. The central and north-western portions of the continent, comprising 46 per cent. of the total land mass, experience evaporation more than twice as great as their rainfall; it is noteworthy that the vegetation over most of this region is characterised by acacia, semi-desert, shrub steppe and porcupine grass. Since the loss by evaporation depends largely on the exposed area, tanks and dams so designed that the surface shall be a minimum are advantageous. Further, the more they are protected from the direct rays of the sun and from winds by means of suitable tree planting, the less will be the loss by evaporation. These matters are naturally of more than ordinary concern in the drier districts of Australia.

(ii) *Comments on Map of Average Annual Evaporation.* The map of average annual evaporation in Australia (see page 36) has been compiled on the basis of records obtained from a number of evaporimeters supplemented by estimates derived from records of saturation deficit by applying the Waite Institute factor of 263.* Some modification of the latter values was found to be necessary in comparison with recordings of evaporimeters.

The standard evaporation tank used in Australia is cylindrical in form and is 36 inches in diameter and 36 inches deep. It is surrounded by a 6-inch water jacket and the whole is sunk into the ground so that the water surface is approximately at ground level.

* Prescott, J. A. "Atmospheric Saturation Deficit in Australia" (Trans. Royal Society, S.A. Vol. LV., 1931).

Saturation deficit is obtained from readings of dry and wet bulb thermometers exposed in a standard Stevenson thermometer shelter. Saturation deficit is the difference between the vapour pressure indicated by the dry and wet bulb readings, and the saturation vapour pressure corresponding to the dry bulb temperature.

The Waite formula, $e = 263 \text{ s.d.}$, is not an exact relationship, but it takes account of one of the major factors in evaporation, i.e., the difference between saturation vapour pressures at the mean dew point and at the mean air temperature. Errors in the formula are found to be fairly consistent in considerable areas of Australia and corrections have been applied accordingly. No evaporation records are available north of latitude 20° , and corrections have been extrapolated for these areas. The evaporation stations on which estimates for the tropics have been based are Alice Springs (N.T.) and Winton (Q'land), and to a lesser degree Blackall (Q'land) and Marble Bar (W.A.).

The map thus presents an estimate of evaporation for which allowance should be made for a certain margin of error (perhaps 10 per cent. or so) on the conservative side. In the absence of definite information, such a map should serve a useful purpose as a basis for many climatic studies.

For graphs and tables of mean monthly evaporation and rainfall at certain selected stations see Official Year Book No. 37, pp. 34-35.

5. **Rainfall.**—(i) *General.* The rainfall of any region is determined mainly by the direction and route of the prevailing winds, by the varying temperatures of the earth's surface over which they blow, and by its physiographical features.

Australia lies within the zones of the south-east trades and "prevailing" westerly winds. The southern limit of the south-east trade strikes the eastern shores at about 30° south latitude, and, with very few exceptions, the heaviest rains of the Australian continent are precipitated along the Pacific slopes to the north of that latitude, the varying quantities being more or less regulated by the differences in elevation of the shores and of the chain of mountains from the New South Wales northern border to Thursday Island, upon which the rain-laden winds blow. The converse effect is exemplified on the north-west coast of Western Australia, where the prevailing winds, blowing from the interior of the continent instead of from the ocean, result in the lightest coastal rain in Australia.

The westerly winds, which skirt the southern shores, are responsible for the reliable, generally light to moderate rains enjoyed by the south-western portion of Western Australia, the agricultural areas of South Australia, a great part of Victoria, and the whole of Tasmania.

(ii) *Distribution of Rainfall.* The average annual rainfall map of Australia (page 37) shows that the heaviest yearly falls occur on the north coast of Queensland (up to more than 160 inches) and in Western Tasmania (up to 140 inches), while from 50 to over 60 inches are received on parts of the eastern seaboard from Jervis Bay (New South Wales) to the northern part of Cape York Peninsula, also around Darwin (Northern Territory), on the West Kimberley coast, near Cape Leeuwin (Western Australia), about the Australian Alps in eastern Victoria and New South Wales, and on the north-eastern highlands in Tasmania. A great part of the interior of the continent, stretching from the far west of New South Wales and the south-west of Queensland to the vicinity of Shark Bay in Western Australia, has a very low average rainfall of less than 10 inches a year. Between these two regions of heavy and very low rainfall are the extensive areas which experience useful to good rains, and in the southern and eastern parts of which are found the best country and most of the population and primary production.

(iii) *Factors Determining Occurrence, Intensity and Seasonal Distribution of Rainfall.* Reference has already been made to the frequent rains occurring in the north-eastern coastal districts of Queensland with the prevailing south-east trade winds and to similar rains in the west of Tasmania with the prevailing westerly winds. Other rains in Australia are associated mainly with tropical and southern depressions.

The former chiefly affect the northern, eastern, and to some extent the central parts of the continent and operate in an irregular manner during the warmer half of the year, but principally from December to March. They vary considerably in activity and scope from year to year, occasionally developing into severe storms off the east and north-west

coasts. Tropical rainstorms sometimes cover an extensive area, half of the continent on occasions receiving moderate to very heavy falls during a period of a few days. Rain is also experienced, with some regularity, with thunderstorms in tropical areas, especially near the coast. All these tropical rains, however, favour mostly the northern and eastern parts of the area referred to; the other parts further inland receive lighter, less frequent and less reliable rainfall. With the exception of districts near the east coast, where some rain falls in all seasons, the tropical parts of the continent receive useful rains only on rare occasions from May to September.

The southern depressions are most active in the winter—June to August—and early spring months. The rains associated with them are fairly reliable and frequent over Southern Australia and Tasmania, and provide during that period the principal factor in the successful growing of wheat. These depressions also operate with varying activity during the remainder of the year, but the accompanying rains are usually lighter. The southern rains favour chiefly the south-west of Western Australia, the agricultural districts of South Australia, Victoria, Tasmania, and the southern parts of New South Wales. They sometimes extend into the drier regions of the interior, but only infrequently and irregularly.

The map showing mean monthly distribution of rainfall over Australia (page 38) gives in graphic form information on the amount and occurrence of rain.

(iv) *Wettest and Driest Regions.* The wettest known part of Australia is on the north-east coast of Queensland, between Port Douglas and Cardwell, where Deeral on the north coast-line has an average annual rainfall of 175.96 inches and Tully on the Tully River 187.19 inches. In addition, three stations situated on, or adjacent to, the Johnstone and Russell Rivers have an average annual rainfall of between 144 and 169 inches. The maximum and minimum annual amounts there are:—Deeral, 287.18 in 1945 and 99.60 inches in 1947, or a range of 187.58 inches; Tully, 310.92 in 1950 and 104.98 inches in 1943, or a range of 205.94 inches; Goondi, 241.53 in 1894 and 67.88 inches in 1915, or a range of 173.65 inches; Innisfail, 232.06 in 1950 and 69.87 inches in 1902, or a range of 162.19 inches; Harvey Creek, 254.77 in 1921 and 80.47 inches in 1902, or a range of 174.30 inches.

On five occasions more than 200 inches have been recorded at Goondi, the last of these being in 1950, when 204.97 inches were registered. The records at this station cover a period of 64 years.

In twenty-four years of record Tully has exceeded 200 inches on ten occasions, whilst in a record of 28 complete years Harvey Creek has four times exceeded this figure.

In Tasmania the wettest part is in the West Coast region, the average annual rainfall at Lake Margaret being 145.53 inches, with a maximum of 177.30 inches in 1948.

The driest known part of the continent is in an area of approximately 180,000 square miles surrounding Lake Eyre in South Australia, where the annual average is between 4 and 6 inches and where the fall rarely exceeds 10 inches for 12 months.

Records of stations have at times been interrupted, but of the 23 stations in this region which have an annual average of less than 5 inches, six have complete records extending from 30 to 55 years. Of these Mulka has the lowest average of 4.13 inches (33 years), followed by Troudaninna with an average of 4.15 inches in 42 years. Troudaninna in the period 1893 to 1936 had only one year in which the total exceeded 9 inches (11.07 inches in 1894). There have been protracted periods when the average has even been less than 3 inches. From 1895 to 1903 Troudaninna received the following annual totals:—2.78, 0.99, 5.71, 3.04, 3.18, 2.83, 1.80, 1.11, 4.87, an average of 2.91 inches. From 1918 to 1929 the average was only 2.65 inches, and in this period from December, 1924 to November, 1929 the average was only 1.70 inches.

Mulka since 1918 has only once exceeded 10 inches for the annual total (11.72 inches in 1920), and in 33 years on 15 occasions the annual total has been less than 3 inches. In one particular period from October, 1926 to September, 1930, the average was only 1.26 inches (505 points in 48 months). However, at Kanowana, an even lower four-year average of 1.12 inches was recorded between 1896 and 1899 with yearly totals of 43, 225, 87 and 94 points. An even smaller total than 43 points was recorded at Mungeranie in 1889 when only 39 points was recorded on 5 days.

The average number of rain days per month in this region is only 1-2 and the annual number ranges between 10 and 20. Oodnadatta (standard 30 years' average rainfall equal to 4.44 inches) has an average of 20 days of rain per year while Cordillo Downs in the extreme north-east corner of the State of South Australia receives 5.16 inches on 12 days per year, averaging about one day of rain each month in the thirty years' period 1911-1940.

No part of the earth, so far as is known, is absolutely rainless, and although at Arica, in northern Chile, the rainfall over a period of 15 years was nil, a further two years in which there were three measurable showers made the "average" for 17 years 0.02 inches.

(v) *Quantities and Distribution of Rainfall.* The general distribution is best seen from the rainfall map (page 37) which shows the areas subject to average annual rainfalls lying between certain limits. The proportions of the total area of each State and of Australia as a whole enjoying varying quantities of rainfall determined from the latest available information are shown in the following table:—

AVERAGE ANNUAL RAINFALL DISTRIBUTION.
(Per Cent.)

Average Annual Rainfall.	N.S.W. (a)	Victoria.	Queens-land.	South Australia	Western Australia.	Tasmania. (b)	Northern Territory	Total.
Under 10 inches ..	19.7	Nil	13.0	82.8	58.0	Nil	24.7	37.6
10—15 " ..	23.5	22.4	14.4	9.4	22.4	Nil	32.4	19.9
15—20 " ..	17.5	15.2	19.7	4.5	6.8	0.7	9.7	10.9
20—25 " ..	14.2	17.9	18.8	2.2	3.7	11.0	6.6	9.1
25—30 " ..	9.1	18.0	11.6	0.8	3.7	11.4	9.3	7.3
30—40 " ..	9.9	16.1	11.1	0.3	3.3	20.4	4.7	6.6
Over 40 " ..	6.1	10.4	11.4	Nil	2.1	56.5	12.6	8.6
Total ..	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

(a) Includes Australian Capital Territory.
available.

(b) Over an area of 2,777 square miles no records are

Referring first to the capital cities, the records of which are given in the next table, it will be seen that Sydney, with an average rainfall of 46.86 inches, occupies the chief place: Brisbane, Perth, Melbourne, Hobart, Canberra and Adelaide follow in that order, Adelaide with 21.03 inches being the driest. The extreme range from the wettest to the driest year is greatest at Brisbane (72.09 inches) and least at Adelaide (19.46 inches).

In order to show how the rainfall is distributed throughout the year in various parts of the continent, average figures for the various climatological districts have been selected. (See map on p. 38). The figures for Northern Rivers (District 14), show that nearly the whole of the rainfall occurs there in the summer months, while little or none falls in the middle of the year. The figures for the Central Coast, south-west of Western Australia (District 9), are the reverse, for while the summer months are dry, the winter months are very wet. In the districts containing Melbourne and Hobart the rain is fairly well distributed throughout the twelve months, with a maximum in October for both districts. In Queensland, the heaviest rains fall in the summer months, but good averages are also maintained during the other seasons in eastern parts.

On the coast of New South Wales, the first half of the year is the wettest, with heaviest falls in the autumn; the averages during the last six months are fair, and moderately uniform. Generally it may be said that approximately one-third of the area of the continent, principally in the eastern and northern parts, enjoys an annual average rainfall of from 20 to 50 inches or more, the remaining two-thirds averaging from 5 to 20 inches.

(vi) *Tables of Rainfall.* The table of rainfall for a fairly long period of years for each of the various Australian capitals affords information as to the variability of the fall in successive years, and the list of the more remarkable falls furnishes information as to what may be expected on particular occasions.

RAINFALL : AUSTRALIAN CAPITAL CITIES.

Year.	CANNBERRA.(a)		PERTH.		ADELAIDE.		BRISBANE.		SYDNEY.		MELBOURNE.		HOBART.(b)	
	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.
	£.		in.		in.		in.		in.		in.		in.	
1920	40.35	124	26.70	119	39.72	122	43.42	159	28.27	162	18.00	182
21	41.09	135	22.64	100	54.31	167	43.34	140	29.76	154	18.04	159
22	31.86	135	23.20	117	35.82	109	39.35	136	25.02	151	28.27	189
23	44.47	134	29.79	139	23.27	93	37.01	123	22.64	158	32.93	198
24	33.79	119	23.44	143	41.08	114	37.01	136	36.48	171	28.76	197
25	31.41	126	21.91	118	53.10	139	50.35	145	17.57	144	22.67	170
26	49.22	167	22.20	116	30.82	111	37.07	127	20.51	149	25.79	187
27	36.59	133	16.92	101	62.08	130	48.56	138	17.98	135	20.13	185
28 ..	18.59	90	44.88	140	19.43	107	52.64	145	40.07	130	24.09	151	30.23	205
29 ..	23.12	70	36.77	132	17.51	119	39.78	118	57.90	129	28.81	168	26.55	194
30 ..	17.33	82	39.80	129	18.65	116	41.22	144	44.47	141	25.41	145	19.38	152
31 ..	24.02	103	39.18	118	22.26	145	66.72	136	49.22	153	28.63	164	27.17	179
32 ..	20.18	118	39.40	121	25.04	141	24.79	97	37.47	146	31.08	179	30.29	155
33 ..	20.78	96	32.47	116	22.12	130	49.71	118	42.71	153	22.28	136	23.18	182
34 ..	35.58	131	40.61	120	20.24	125	54.26	117	64.91	183	33.53	157	23.17	194
35 ..	23.78	95	32.28	129	23.45	140	34.64	111	30.97	131	29.98	183	32.22	196
36 ..	26.24	108	30.64	118	19.34	121	21.77	101	30.22	130	24.30	187	19.60	178
37 ..	20.46	82	35.28	120	23.01	128	34.79	113	52.00	157	21.45	144	20.65	160
38 ..	19.26	79	29.64	111	19.26	119	43.49	110	39.17	132	17.63	131	31.32	169
39 ..	27.63	116	45.70	123	23.29	139	41.43	122	33.67	127	33.11	166	27.23	188
40 ..	17.38	64	20.00	98	16.16	116	42.37	93	39.34	125	19.83	126	17.17	135
41 ..	19.55	91	34.74	122	22.56	126	31.50	105	26.74	129	31.78	157	23.49	145
42 ..	25.76	104	39.24	140	25.44	133	44.01	125	48.29	121	29.79	148	19.42	163
43 ..	24.59	123	31.46	117	17.84	135	50.68	126	50.74	136	18.80	150	20.84	149
44 ..	12.05	75	27.39	123	17.13	114	27.85	100	31.04	115	21.32	143	26.23	151
45 ..	22.35	100	52.67	137	17.85	105	48.16	130	46.47	136	19.22	152	16.92	157
46 ..	22.31	94	41.47	122	22.59	135	38.66	83	36.05	111	29.80	177	39.45	193
47 ..	27.95	135	43.42	137	21.89	146	60.30	146	41.45	137	30.47	163	38.61	181
48 ..	32.11	101	34.75	126	21.40	122	41.54	106	38.83	131	20.98	155	23.42	178
49 ..	27.71	100	27.15	126	18.23	119	47.18	121	66.26	149	31.41	163	22.85	157
50 ..	43.35	132	32.27	122	16.06	91	63.93	152	86.33	183	26.18	147	19.25	131
Average No. of Years Standard 30 years' Normal	24.00	100	34.91	122	21.03	123	44.93	125	46.86	151	25.62	143	24.51	168
	23	23	75	75	112	112	99	91	92	92	95	95	68	68
	35.99	128	21.09	122	40.09	117	44.80	143	25.89	156	25.03	180

(a) Commonwealth Forestry Bureau; records in issues prior to No. 36 were for the station at Acton which closed down in 1939. (b) Records taken from present site commenced 1883.

6. Remarkable Falls of Rain.—The following are the most notable falls of rain in the various States and Territories which have occurred within a period of twenty-four hours. For other very heavy falls at various localities reference may be made to Official Year Book No. 14, pp. 60-64, No. 22, pp. 46-48 and No. 29, pp. 43, 44 and 51 :—

HEAVY RAINFALLS : NEW SOUTH WALES, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Ammt.		Name of Town or Locality.	Date.	Ammt.	
		in.				in.	
Bega ..	27 Feb., 1919	17.88		Madden's Creek	13 Jan., 1911	18.68	
Broger's Creek ..	14 Feb., 1898	20.05		Morpeth ..	9 Mar., 1893	21.52	
" ..	13 Jan., 1911	20.83		Mt. Kembla ..	13 Jan., 1911	18.25	
Buladelah ..	16 Apr., 1927	19.80		Mt. Pleasant ..	5 May, 1925	20.10	
Candelo ..	27 Feb., 1919	18.58		Nimbin ..	6 Feb., 1939	16.26	
Condong ..	27 Mar., 1887	18.66		South Head (Sydney Harbour)	16 Oct., 1844	20.41	
Cordeaux River..	14 Feb., 1898	22.58		" ..	29 Apr., 1841	20.12	
Dorrigo ..	24 June, 1950	25.04		" ..	5 Mar., 1893	20.00	
Foxground ..	11 Sept., 1950	17.04		Towamba ..	15 Mar., 1936	20.00	
Kembla Heights	13 Jan., 1911	17.46		Viaduct Creek ..			

HEAVY RAINFALLS : QUEENSLAND, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		in.			in.
Babinda (Cairns) ..	2 Mar., 1935	24.14	Kuranda (Cairns) ..	2 Apr., 1911	28.80
Banyan (Cardwell)	12 Feb., 1927	24.00	Landsborough ..	2 Feb., 1893	25.15
Buderim Mountain	11 Jan., 1898	26.20	Macnade Mill ..	6 Feb., 1901	23.33
Carruchan ..	24 Jan., 1934	24.00	Plane Creek (Mackay)	26 Feb., 1913	27.73
Crohamhurst			Port Douglas ..	1 Apr., 1911	31.53
(Blackall Range)	2 Feb., 1893	35.71	Sarina ..	26 Feb., 1913	27.75
Deeral ..	2 Mar., 1935	27.60	Springbrook ..	24 Jan., 1947	27.07
Flat Top Island ..	21 Jan., 1918	25.18	Tully Mill ..	12 Feb., 1927	23.86
Goondi ..	30 Jan., 1913	24.10	Woodlands (Yepp'n)	3 Jan., 1893	23.07
Harvey Creek ..	3 Jan., 1911	27.75	Yarrabah ..	2 Apr., 1911	30.65

HEAVY RAINFALLS : WESTERN AUSTRALIA, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		in.			in.
Balla Balla ..	21 Mar., 1899	14.40	Pilbara ..	2 Apr., 1898	14.04
Boodarie ..	21 Mar., 1899	14.53	Roebuck Plains ..	5 Jan., 1917	14.01
Broome ..	6 Jan., 1917	14.00	" "	6 Jan., 1917	22.36
Carlton Hill ..	7 Feb., 1942	12.75	Thangoo ..	17-19 Feb. '96	24.18
Derby ..	7 Jan., 1917	16.47	Towrana ..	1 Mar., 1943	12.16
Fortesque ..	3 May, 1890	23.36	Whim Creek ..	3 Apr., 1898	29.41
Jimba Jimba ..	1 Mar., 1943	11.54	Winderrrie ..	17 Jan., 1923	14.23
Marble Bar ..	2 Mar., 1941	12.00	Widjip ..	1 Apr., 1934	19.54

HEAVY RAINFALLS : NORTHERN TERRITORY, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		in.			in.
Bathurst Island			Cape Don ..	13 Jan., 1935	13.58
Mission ..	7 Apr., 1925	11.85	Darwin ..	7 Jan., 1897	11.67
Borrooloola ..	14 Mar., 1899	14.00	Groote Eylandt ..	9 Apr., 1931	14.29
Brock's Creek ..	24 Dec., 1915	14.33	Timber Creek ..	5 Feb., 1942	13.65

HEAVY RAINFALLS : SOUTH AUSTRALIA, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		in.			in.
Ardrrossan ..	18 Feb., 1946	8.10	Mannum ..	25 Jan., 1941	6.84
Cape Willoughby ..	18 Feb., 1946	6.80	Port Victoria ..	18 Feb., 1946	7.08
Carpa ..	18 Feb., 1946	7.83	Torrens Vale ..	25 Jan., 1941	6.77
Edithburg ..	18 Feb., 1946	7.46	Wilmington ..	1 Mar., 1921	7.12
Hesso ..	18 Feb., 1946	7.36	Wirrabara ..	7 Mar., 1910	6.80
Maitland ..	18 Feb., 1946	7.21	Wynbring ..	28 Feb., 1921	7.70

HEAVY RAINFALLS : VICTORIA, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		in.			in.
Blackwood "Green-hill" ..	26 Jan., 1941	8.98	Kalorama ..	1 Dec., 1934	10.05
Cann River ..	27 Feb., 1919	9.56	Korumburra ..	1 Dec., 1934	8.51
" "	16 Mar., 1938	9.94	Mt. Buffalo ..	6 June, 1917	8.53
Corinella ..	28 June, 1943	8.75	Olinda ..	1 Dec., 1934	9.10
Erica ..	1 Dec., 1934	8.66	Tambo Crossing ..	13 July, 1925	8.89
Hazel Park ..	1 Dec., 1934	10.50	Tonghi Creek ..	27 Feb., 1919	9.90

HEAVY RAINFALLS : TASMANIA, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		in.			in.
Cullenswood ..	5 Apr., 1929	11.12	Riana ..	5 Apr., 1929	11.08
Gould's Country ..	8-10 Mar., '11	15.33	The Springs ..	30-31 Jan., '16	10.75
Lottah ..	8-10 Mar., '11	18.10	Triabunna ..	5 June, 1923	10.20
Mathinna ..	5 Apr., 1929	13.25			

HEAVY RAINFALLS : AUSTRALIAN CAPITAL TERRITORY, UP TO 1950, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		in.			in.
Canberra (Acton) ..	27 May, 1925	6.84	Land's End ..	27 May, 1925	6.35
Cotter Junction ..	27 May, 1925	7.13	Uriarra (Woodside)	27 May, 1925	6.57

7. **Snowfall.**—Light snow has been known to fall occasionally as far north as latitude 31° S., and from the western to the eastern shores of the continent. During exceptional seasons, it has fallen simultaneously over two-thirds of New South Wales, and has extended at times along the whole of the Great Dividing Range, from its southern extremity in Victoria as far north as Toowoomba in Queensland. During the winter, for several months, snow covers the ground to a great extent on the Australian Alps, where also the temperature falls below zero Fahrenheit during the night. In the ravines around Mt. Kosciusko and similar localities the snow never entirely disappears after a severe winter.

8. **Hail.**—Hail falls most frequently along the southern shores of the continent in the winter, and over eastern Australia during the summer months. The size of the hailstones generally increases with distance from the coast. A summer rarely passes without some station experiencing a fall of stones exceeding in size an ordinary hen-egg, and many riddled sheets of light-gauge galvanized iron bear evidence of the weight and penetrating power of the stones.

The hailstones occur most frequently when the barometric readings indicate a flat and unstable condition of pressure. Tornadoes or tornadic tendencies are almost invariably accompanied by hail, and on the east coast the clouds from which the stones fall are frequently of a remarkable sepia-coloured tint.

9. **Barometric Pressures.**—The mean annual barometric pressure (corrected to sea level and standard gravity) in Australia varies from 29.80 inches on the north coast to 29.92 inches over the central and 30.03 inches in the southern parts of the continent. In January, the mean pressure ranges from 29.70 inches in the northern and central areas to 29.95 inches in the southern. The July mean pressure ranges from 29.90 inches at Darwin to 30.12 inches at Alice Springs. Barometer readings corrected to mean sea level and standard gravity have, under anticyclonic conditions, ranged as high as 30.935 inches (at Hobart on 13th July, 1846) and have fallen as low as 27.55 inches. This lowest record was registered at Mackay during a tropical hurricane on 21st January, 1918. An almost equally abnormal reading of 27.88 inches was recorded at Innisfail during a similar storm on 10th March, 1918. For graphs of Mean Barometric Pressure at Capital Cities see Official Year Book No. 37, p. 35.

10. **Wind.**—(i) *Trade Winds.* The two distinctive wind currents in Australia are, as previously stated, the south-east trade and the "prevailing" westerly winds. As the belt of the earth's atmosphere in which they blow apparently follows the sun's ecliptic path north and south of the equator, so the area of the continent affected by these winds varies at different seasons of the year. During the summer months the anticyclonic belt travels in high latitudes, thereby bringing the south-east trade winds as far south as 30° south latitude. The "prevailing" westerly winds retreat a considerable distance to the south of Australia, and are less in evidence in the hot months. When the sun passes to the north of the equator, the south-east trade winds follow it, and only operate to the north of the tropics for the greater part of the winter. The westerly winds come into lower latitudes during the same period of the year. They sweep across the southern areas of the continent from Cape Leeuwin to Cape Howe, and during some seasons are remarkably persistent and strong, and occasionally penetrate to almost tropical latitudes.

(ii) *North-west Monsoon.* As the belt of south-east trade winds retreats southward during the summer, it is replaced in the north and north-west of Australia first by a sequence of light variable winds and then by the north-west monsoon. In Australia, the north-west monsoon has not the persistence nor regularity of the Indian south-west

monsoon but is sufficiently characteristic for the summer in the north of Australia to be called the "North-west Season". In central and eastern Queensland, the north-west monsoon in the summer has comparatively little effect and the trade winds, albeit weakened, are still dominant winds. With the migration of the sun northward in the autumn, the north-west monsoon is itself replaced first by light variable winds and then by the trade winds.

Further particulars of Australian wind conditions and meteorology will be found in Official Year Book No. 38, pp. 58-61.

(iii) *Cyclones and Storms.* The "elements" in Australia are ordinarily peaceful, and while destructive cyclones have visited various parts, more especially coastal areas, such visitations are rare, and may be properly described as erratic.

During the winter months, the southern shores of the continent are subject to deep depressions of the southern low-pressure belt. They are felt most severely over the south-western parts of Western Australia, to the south-east of South Australia, in Bass Strait, including the coast-line of Victoria, and on the west coast of Tasmania. Apparently the more violent wind pressures from these disturbances are experienced in their northern half, or in that part of them which has a north-westerly to a south-westerly circulation.

The north-east coast of Queensland is occasionally visited by hurricanes from the north-east tropics. During the first four months of the year, these hurricanes appear to have their origin in the neighbourhood of the South Pacific Islands, their path being a parabolic curve first to the south-west and finally towards the south-east.

Very severe cyclones, locally known as "willy willies," are peculiar to the north-west coast of Western Australia from the months of November to April, inclusive. They usually originate over the ocean to the north or north-west of Australia, and travel in a south-westerly direction with continually increasing force, displaying their greatest energy near Cossack and Onslow, between latitudes 20° and 22° South. The winds in these storms, like those from the north-east tropics, are very violent and destructive. The greatest velocities are usually to be found in the south-eastern quadrant of the cyclones, with north-east to east winds. After leaving the north-west coast, these storms either travel southwards, following the coast-line, or cross the continent to the Great Australian Bight. When they take the latter course, their track is marked by torrential rains, as much as 29.41 inches, for example, being recorded in 24 hours at Whim Creek from one such occurrence. Falls of 10 inches and over have frequently been recorded in the northern interior of Western Australia from similar storms.

Some further notes on severe cyclones and on "southerly bursters", a characteristic feature of the eastern part of Australia, appear in previous issues of the Official Year Book (see No. 6, pp. 84-86), and a special article dealing with "Australian Hurricanes and Related Storms" appears in Official Year Book No. 16, pp. 80-84.

Depressions vary considerably in their isobaric forms, intensity and other characteristics. Some bring rain in variable quantities, some heat and others mainly wind. A common type in southern Australia is the "Δ" shaped trough with an abrupt "backing" of the wind or "line squall" as it passes. The cold front is most frequently found through the centre of the "trough" because it is along this line, and extending into the upper levels of the atmosphere that the demarcation of different air masses is so well defined. The best rains occur in inland Australia when extensive masses of warm moist tropical air move into the interior and are forced to rise by convergence of flow or by impact with a cold air stream.

The speed of low pressure systems is very variable, but in general in southern latitudes the movement is of the order of 500 to 700 miles per day.

11. **Influences affecting Australian Climate.**—(i) *General.* Australian history does not cover a sufficient period, nor is the country sufficiently occupied, to ascertain whether or not the advance of settlement has materially affected the climate as a whole. Local changes have, however, taken place, a fact which suggests that settlement and the treatment of the land have a distinct effect on local conditions. For example, low-lying lands on the north coast of New South Wales, which originally were seldom subject to frosts, have, with the denudation of the surrounding hills from forests, experienced annual visitations, the probable explanation being that through the absence of trees the cold air of the highlands now flows unchecked and untempered down the sides of the hills to the valleys and lower lands.

(ii) *Influence of Forest on Climate.* As already indicated, forests doubtless exercise a great influence on local climate, and hence, to the extent that forestal undertakings will allow, the weather can be controlled by human agency. The direct action of forests is an equalizing one; thus, especially in equatorial regions, and during the warmest portion of the year, they considerably reduce the mean temperature of the air. They also reduce the diurnal extremes of shade temperatures by altering the extent of radiating surface by evaporation, and by checking the movement of air, and while decreasing evaporation from the ground, they increase the relative humidity. Vegetation greatly diminishes the rate of flow-off of rain and the washing away of surface soil, and when a region is protected by trees a steadier water supply is ensured, and the rainfall is better conserved. In regions of snowfall, the supply of water to rivers is similarly regulated, and without this and the sheltering influence of ravines and "gullies," watercourses supplied mainly by melting snow would be subject to alternative periods of flooding and dryness. This is borne out in the case of the inland rivers, the River Murray, for example, which has never been known to become dry, deriving its steadiness of flow mainly through the causes indicated.

(iii) *Direct Influence of Forests on Rainfall.* Whether forests have a direct influence on rainfall is a debatable question, some authorities alleging that precipitation is undoubtedly induced by forests, while others take the opposite view.

Sufficient evidence exists, however, to prove that, even if the rainfall has not increased, the beneficial climatic effect of forest lands more than warrants their protection and extension. Rapid rate of evaporation, induced by both hot and cold winds, injures crops and makes life uncomfortable on the plains, and, while it may be doubted that the forest aids in increasing precipitation, it must be admitted that it does check winds and the rapid evaporation due to them. Trees as wind-breaks have been successfully planted in central parts of the United States of America, and there is no reason why similar experiments should not be successful in many parts of the treeless interior of Australia. The belts should be planted at right angles to the direction of the prevailing parching winds, and if not more than half a mile apart will afford shelter to the enclosed areas.

12. **Rainfall and Temperatures, Various Cities.**—The Official Year Book No. 34, p. 28, shows rainfall and temperature for various important cities throughout the world, and for the Australian capitals.

13. **Climatological Tables.**—The averages and extremes for a number of climatological elements, which have been determined from long series of observations at the Australian capitals up to and including the year 1950, are given on pp. 58–64.

NOTE.—The following points apply throughout:—

- (i) Where records are available, mean or average values have been calculated on a standard period of 30 years from 1911 to 1940.
- (ii) Extreme values have been extracted from all available years of actual record, but the number of years quoted does not include intervening periods when observations were temporarily discontinued.

CLIMATOLOGICAL DATA : CANBERRA, AUSTRALIAN CAPITAL TERRITORY.

LAT. 35° 18' S., LONG. 149° 06' E. HEIGHT ABOVE M.S.L. 1,906 FT.

BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS AND CLEAR DAYS.

Month.	Bar. corrected to 32° F. Mm. Sea level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 20 feet.)				Prevailing Direction.	Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m. and 3 p.m.(a)	No. of Clear Days.	
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevaling Direction.						
					9 a.m.						3 p.m.
No. of years of observations.	20	21	22	(b)	23	23	22	14	20	21	
January ..	29.837	5.1	14.9 23/33	—	NW	NW	8.73	1.2	4.7	7.7	
February ..	29.899	4.5	15.2 24/33	—	E	NW	6.23	3.0	4.8	6.3	
March ..	30.012	4.1	18.2 28/42	—	E	NW	5.40	0.1	5.0	6.7	
April ..	30.076	3.8	18.6 8/45	—	NW	NW	3.36	0.3	5.4	4.7	
May ..	30.151	3.1	12.6 3/30	—	NW	NW	2.03	0.1	5.4	5.7	
June ..	30.141	3.7	16.1 2/30	—	NW	NW	1.32	0.1	6.1	4.1	
July ..	30.129	3.6	23.4 7/31	—	NW	NW	1.30	0.0	5.6	5.0	
August ..	30.077	4.3	15.7 25/36	—	NW	NW	1.86	0.1	5.4	5.0	
September ..	30.043	4.7	17.4 38/34	—	NW	NW	3.01	0.4	5.0	6.1	
October ..	29.960	4.6	12.4 27/40	—	NW	NW	4.64	1.0	5.2	5.4	
November ..	29.897	4.9	17.2 28/42	—	NW	NW	6.02	1.2	5.5	4.2	
December ..	29.837	5.0	16.1 11/38	—	NW	NW	7.92	1.0	5.1	5.0	
Year { Totals ..	—	—	—	—	—	—	52.42	8.5	—	—	
{ Averages	30.005	4.3	—	—	NW	NW	—	—	5.3	—	
{ Extremes	—	—	23.4 7/7/31	—	—	—	—	—	—	—	

(a) Scale 0-10. (b) No record.

TEMPERATURE AND SUNSHINE.

Month.	Mean Temperature (°Fahr.)			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.	
No. of years over which observation extends.	23	23	23	23	23	23	(a)	23	20
January ..	82.5	56.0	69.3	107.4 11/39	39.4 18/49	68.0	—	30.1 10/50	8.1
February ..	81.1	56.0	68.6	99.8 13/33	35.0 (b)	64.8	—	26.5 23/43	7.5
March ..	75.9	52.4	64.1	99.1 6/38	31.8 31/49	64.3	—	26.4 26/35	7.1
April ..	66.5	45.1	55.8	89.7 6/38	29.0 29/34	60.7	—	19.0 18/44	6.6
May ..	59.5	38.8	49.2	72.6 1/36	22.5 9/29	50.1	—	15.6 (c)	5.2
June ..	52.4	35.4	43.9	61.0 (d)	18.1 20/35	42.9	—	8.9 25/44	4.3
July ..	51.7	33.9	42.8	63.5 16/34	20.0 (e)	43.5	—	10.8 9/37	4.7
August ..	55.2	35.4	45.3	70.5 28/34	21.0 3/29	49.5	—	10.7 6/44	5.0
September ..	61.3	39.0	50.1	81.5 16/34	25.2 6/46	56.3	—	13.0 6/45	7.2
October ..	67.4	44.2	55.8	90.0 13/46	29.0 24/28	61.0	—	18.2 2/45	7.7
November ..	73.4	49.3	61.4	101.4 19/44	32.2 11/36	69.2	—	25.9 6/40	7.9
December ..	79.7	53.6	66.6	103.5 27/38	36.0 24/28	67.5	—	30.2 (f)	8.2
Year { Averages	67.2	44.9	56.1	—	—	—	—	—	6.7
{ Extremes	—	—	—	107.4 11/1/39	18.1 20/6/35	89.3	—	8.9 25/6/44	—

(a) No record. (b) 22/31 and 23/31. (c) 13/37 and 15/46. (d) 3/27 and 28/30. (e) 19/29, 9/37, and 27/43. (f) 2/39 and 20/48.

HUMIDITY, RAINFALL AND FOG.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (Inches).					Fog. Mean No. of Days of Fog.	
		Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.		
											22
No. of years over which observation extends.	22	22	22	22	23	23	23	23	23	23	19
January ..	0.374	52	69	39	2.14	7	6.69 1941	0.02 1932	2.47 19/50	—	0.0
February ..	0.399	57	71	49	2.22	6	6.03 1948	0.01 1933	3.24 17/28	—	0.0
March ..	0.385	66	79	48	2.33	7	12.69 1950	0.01 1940	2.46 19/50	—	0.3
April ..	0.317	71	81	54	2.16	8	3.75 1935	0.07 1942	2.52 9/45	—	1.1
May ..	0.257	79	87	67	1.82	7	6.13 1948	0.06 1935	3.88 3/48	—	4.4
June ..	0.216	82	90	72	1.63	9	6.09 1931	0.18 1944	1.65 24/31	—	6.0
July ..	0.204	81	87	73	1.64	10	4.09 1932	0.27 1940	2.02 13/33	—	5.0
August ..	0.212	75	88	60	1.89	11	4.71 1939	0.36 (a)	2.07 12/29	—	2.0
September ..	0.240	65	74	51	1.59	9	3.03 1937	0.13 1946	1.75 3/47	—	1.0
October ..	0.279	59	72	46	2.53	10	6.59 1934	0.34 1940	2.51 25/34	—	0.2
November ..	0.316	55	67	38	2.15	8	4.45 1950	0.28 1936	2.45 9/50	—	0.0
December ..	0.347	51	70	37	1.90	8	8.80 1947	0.16 1938	2.29 28/29	—	0.0
Year { Totals	—	—	—	—	24.00	100	—	—	—	—	20.0
{ Averages	0.285	66	—	—	—	—	—	—	—	—	—
{ Extremes	—	—	90	37	—	—	12.69 3/50	0.01 2/33/3/40	3.88 3/5/48	—	—

(a) 1944 and 1949.

CLIMATOLOGICAL DATA : PERTH, WESTERN AUSTRALIA.
 LAT. 31° 57' S., LONG. 115° 51' E. HEIGHT ABOVE M.S.L. 210 FT.
 BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS AND CLEAR DAYS.

Month.	Bar. corrected to 32° F. M.S. Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 71 feet.)				Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m. to 3 p.m., 9 p.m. (d)	No. of Clear Days.	
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.					
					9 a.m.					3 p.m.
No. of years of observations.	30(b)	30(b)	52	38	30(b)	30(b)	30(b)	30(b)		
January ..	29.897	13.8	33.2 27/98	49	E	SSW	10.37	2	2.9	14
February ..	29.922	13.5	27.1 6/08	50	ENE	SSW	8.63	2	3.1	13
March ..	29.976	12.8	27.1 6/13	66	E	SSW	7.52	2	3.5	12
April ..	30.071	10.7	39.8 25/00	61	ENE	SSW	4.62	2	4.2	9
May ..	30.062	10.6	34.4 29/32	73	NE	WSW	2.80	3	5.4	6
June ..	30.068	10.6	38.1 17/27	80	N	NW	1.82	2	5.9	5
July ..	30.082	11.2	42.3 20/26	73	NNE	W	1.76	2	5.6	5
August ..	30.084	11.8	40.3 15/03	77	N	WNW	2.37	2	5.6	6
September ..	30.073	11.8	36.0 11/05	75	ENE	SSW	3.44	1	4.9	8
October ..	30.033	12.6	33.7 6/16	63	SE	SW	5.38	1	4.8	8
November ..	29.989	13.4	32.4 18/97	63	E	SW	7.65	2	3.9	9
December ..	29.923	13.9	32.3 6/22	64	E	SSW	9.69	2	3.2	13
Year { Totals ..	—	—	—	—	—	—	66.05	23	—	108
Year { Averages ..	30.015	12.2	—	—	E	SSW	—	—	4.4	—
Year { Extremes ..	—	—	42.3 20/7/26	80	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940).

TEMPERATURE AND SUNSHINE.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.	
No. of years over which observation extends.	30(a)	30(a)	30(a)	54	54	54	52	52	30(a)
January ..	84.6	63.3	73.9	110.2 12/34	48.6 20/25	61.6	177.3 22/14	39.5 20/25	10.4
February ..	85.1	63.5	74.3	112.2 8/33	47.7 1/02	64.5	173.7 4/34	39.8 1/13	9.8
March ..	81.3	61.5	71.4	106.4 14/22	45.8 8/03	60.6	167.0 10/18	36.7 8/03	8.8
April ..	76.3	57.4	66.8	99.7 9/10	39.3 20/14	60.4	157.0 8/16	31.0 20/14	7.5
May ..	69.0	52.8	60.9	90.4 2/07	34.3 11/14	56.1	146.0 4/25	25.3 11/14	5.7
June ..	64.4	49.8	57.1	81.7 2/14	35.0 30/20	46.7	135.5 9/14	26.3 11/37	4.8
July ..	62.8	48.0	55.4	76.4 21/31	34.2 7/16	42.2	133.2 13/15	25.1 30/20	5.4
August ..	63.8	48.4	56.1	82.0 21/40	35.3 31/08	46.7	145.1 29/21	26.7 24/35	6.0
September ..	66.8	50.4	58.6	90.9 30/18	38.5 15/47	52.4	153.6 29/16	29.2 21/16	7.1
October ..	69.7	52.6	61.1	95.3 30/22	40.0 16/31	55.3	157.5 31/36	29.8 16/31	8.2
November ..	76.7	57.3	67.0	104.6 24/13	42.0 1/04	62.6	167.0 30/25	35.5 (b)	9.6
December ..	81.2	60.9	71.0	107.9 20/09	48.0 2/10	59.9	168.8 11/27	39.0 12/20	10.4
Year { Averages ..	73.5	55.5	64.5	—	—	—	177.3 22/1/14	25.1 30/7/20	—
Year { Extremes ..	—	—	—	112.2 8/2/33	34.2 7/7/16	78.0	—	—	7.8

(a) Standard 30 years' normal (1911-1940). (b) 6/10 and 14/12.

HUMIDITY, RAINFALL AND FOG.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).				Fog. Mean No. of Days of Fog.		
		Mean 9 a.m.	Mean.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.		Least Monthly.	Greatest in One Day.
No. of years over which observation extends.	30(a)	30(a)	54	54	30(a)	30(a)	75	75	75	30(a)	
January ..	0.438	51	61	41	0.33	3	2.17 1879	Nil (b)	1.74 27/79	0	
February ..	0.434	51	65	43	0.50	3	2.98 1915	Nil (b)	1.63 26/15	0	
March ..	0.432	57	66	46	0.90	5	5.71 1934	Nil (b)	3.03 9/34	0	
April ..	0.397	61	73	51	1.75	8	5.85 1926	Nil 1920	2.62 30/04	1	
May ..	0.365	70	81	61	5.14	15	12.13 1879	0.98 1903	3.00 17/42	2	
June ..	0.337	75	83	68	7.55	17	18.75 1945	2.16 1877	3.90 10/20	2	
July ..	0.322	76	84	69	7.08	19	12.28 1926	2.42 1876	3.00 4/81	2	
August ..	0.316	71	81	62	5.78	19	12.53 1945	0.46 1902	2.91 14/45	1	
September ..	0.341	66	75	58	3.37	15	7.84 1923	0.34 1916	1.82 4/31	1	
October ..	0.345	60	75	52	2.30	12	7.87 1890	0.15 1946	1.73 3/33	0	
November ..	0.374	52	63	41	0.75	7	2.78 1916	Nil 1891	1.40 15/48	0	
December ..	0.409	51	63	44	0.54	5	3.05 1888	Nil (c)	1.72 1/88	0	
Year { Totals ..	—	—	—	—	35.99	128	—	—	—	8	
Year { Averages ..	0.370	62	—	—	—	—	—	—	—	—	
Year { Extremes ..	—	—	84	41	—	—	18.75 6/1945	Nil Various months	3.90 10/6/20	—	

(a) Standard 30 years' normal (1911-1940). (b) Various years. (c) 1886 and 1924.

CLIMATOLOGICAL DATA : ADELAIDE, SOUTH AUSTRALIA.
 LAT. 34° 56' S., LONG. 138° 35' E. HEIGHT ABOVE M.S.L. 140 FT.
 BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS AND CLEAR DAYS.

Month.	Bar. corrected to 32° F. M.S. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 75 feet.)					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m. to 3 p.m., 9 p.m. (a)	No. of Clear Days
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.					
					9 a.m.	3 p.m.				
No. of years of observations.	30(b)	30(b)	73	34	30(b)	30(b)	30(b)	30(b)	30(b)	
January ..	29.917	9.9	31.6 19/99	72	SW	SW	9.27	2.3	3.6	12.9
February ..	29.953	8.8	28.8 22/96	64	NE	SW	7.56	2.0	3.7	11.2
March ..	30.037	8.3	26.2 9/12	63	S	SW	6.39	1.8	4.0	10.6
April ..	30.119	8.0	32.2 10/96	81	NE	SW	3.78	1.5	5.2	7.2
May ..	30.131	8.1	31.7 9/80	63	NE	NW	2.27	1.3	5.8	4.9
June ..	30.119	8.3	31.3 12/78	67	NE	N	1.37	1.3	6.1	4.1
July ..	30.111	8.5	28.1 25/82	60	NE	NW	1.34	1.5	6.0	4.3
August ..	30.084	9.2	32.2 31/97	57	NE	SW	1.99	2.0	5.5	5.6
September ..	30.050	9.2	30.0 2/87	69	NNE	SW	3.05	2.0	5.3	5.8
October ..	30.007	9.8	32.0 28/98	69	NNE	SW	5.03	2.8	5.3	5.7
November ..	29.990	9.9	32.2 7/48	79	SW	SW	6.68	3.3	4.9	7.2
December ..	29.922	9.9	28.1 12/91	75	SW	SW	8.74	2.2	4.2	9.5
Year { Totals ..	—	—	—	—	—	—	57.68	24.0	—	89.0
Year { Averages	30.037	9.0	—	—	NE	SW	—	—	—	—
Year { Extremes	—	—	32.2 (c)	81	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940). (c) 10/4/1896, 31/8/1897 and 7/11/1948.

TEMPERATURE AND SUNSHINE.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.	
No. of years over which observation extends.	30(a)	30(a)	30(a)	94	94	94	54(b)	90	30(a)
January ..	84.8	61.0	72.9	117.7 12/39	45.1 21/84	72.6	180.0 18/82	36.5 14/79	10.0
February ..	85.7	61.8	73.7	113.6 12/99	45.5 23/18	68.1	170.5 10/00	35.8 23/26	9.3
March ..	81.3	59.1	70.2	110.5 9/34	43.9 21/33	66.6	174.0 17/83	32.1 21/33	7.9
April ..	73.0	54.4	63.7	98.6 5/38	39.6 15/59	59.0	155.0 1/83	30.2 16/17	6.0
May ..	66.8	50.8	58.8	89.5 4/21	36.9 (e)	52.6	148.2 12/79	25.6 19/28	4.8
June ..	61.0	46.6	53.8	76.0 23/65	32.5 (d)	43.5	138.8 18/79	21.0 24/44	4.2
July ..	59.9	45.4	52.7	74.0 11/06	32.0 24/08	42.0	134.5 26/90	22.1 30/29	4.3
August ..	62.3	46.2	54.3	85.0 31/11	32.3 17/59	52.7	140.0 31/92	22.8 11/29	5.4
September ..	66.8	48.3	57.5	91.3 29/44	32.7 4/58	58.6	160.5 23/82	25.0 25/27	6.3
October ..	72.5	51.7	62.1	102.9 21/22	36.0 —/57	66.9	162.0 30/21	27.8 (e)	7.3
November ..	78.1	55.4	66.7	113.5 21/65	40.8 2/09	72.7	166.9 20/78	31.5 2/09	8.6
December ..	82.6	58.9	70.7	114.6 29/31	43.0 (f)	71.6	175.7 7/99	32.5 4/84	9.5
Year { Averages	72.9	53.3	63.1	—	—	—	180.0	21.0	7.0
Year { Extremes	—	—	—	117.7 12/1/39	32.0 24/7/08	85.7	180.0 18/1/82	24.6/44	—

(a) Standard 30 years' normal (1911-1940). (b) Records incomplete, 1931-34. Discontinued, 1934.
 (c) 26/1895. (d) 27/1876 and 24/1944. (e) 4/1931 and 2/1918. (f) 16/1861 and 4/1906.

HUMIDITY, RAINFALL AND FOG.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) 9 a.m.			Rainfall (inches).				Fog. Mean No. of Days of Fog		
		Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.		Greatest in One Day.	
											30(a)
No. of years over which observation extends.	30(a)	30(a)	83	83	30(a)	30(a)	112	112	112	30(a)	
January ..	0.327	39	59	29	0.76	5	4.00	1850	Nil (b)	2.30 2/89	0.0
February ..	0.352	41	56	30	1.10	5	6.09	1925	Nil (b)	5.57 7/25	0.0
March ..	0.332	44	58	29	0.87	5	4.60	1878	Nil (b)	3.50 5/78	0.0
April ..	0.329	55	72	37	1.45	10	6.78	1853	Nil	1945 3.15 5/60	0.0
May ..	0.313	64	76	49	2.49	13	7.75	1875	0.10	1934 2.75 1/53	0.6
June ..	0.294	75	84	67	2.93	15	8.58	1916	0.42	1886 2.11 1/20	1.1
July ..	0.282	75	87	66	2.49	16	5.38	1865	0.37	1899 1.75 10/65	1.4
August ..	0.282	68	78	54	2.58	16	6.24	1852	0.33	1944 2.23 19/51	0.4
September ..	0.289	59	72	44	2.39	13	5.83	1923	0.45	1896 1.59 20/23	0.2
October ..	0.287	48	67	29	1.54	10	4.38	1948	0.17	1914 2.24 16/08	0.0
November ..	0.292	41	57	31	1.22	8	4.10	1934	0.04	1885 2.08 7/34	0.0
December ..	0.322	40	50	31	1.27	6	3.98	1861	Nil	1904 2.42 23/13	0.0
Year { Totals ..	—	—	—	—	21.09	122	—	—	—	—	3.7
Year { Averages	0.304	52	—	—	—	—	—	—	—	—	—
Year { Extremes	—	—	87	29	—	—	8.58 6/1916	Nil (c)	5.57 7/2/25	—	—

(a) Standard 30 years' normal (1911-1940). (b) Various years. (c) December to April, various years.

CLIMATOLOGICAL DATA : BRISBANE, QUEENSLAND.

LAT. 27° 28' S., LONG. 153° 2' E. HEIGHT ABOVE M.S.L. 134 FT.
 BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS AND CLEAR DAYS.

Month.	Bar. corrected to 32° F. M.S. Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind.				Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m. (e)	No. of Clear Days.
		(Height of Anemometer 105 feet.)							
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.				
				9 a.m.	3 p.m.				
No. of years of observations.	30(b)	30(b)	36	36	30(b)	30(b)	30(b)	30(b)	30(b)
January ..	29.865	6.8	19.7 23/47	58	SE	NE	6.74	9.8	5.7 3.5
February ..	29.912	7.0	21.0 5/31	67	SE	NE	5.49	6.5	5.6 2.4
March ..	29.975	6.5	20.3 1/29	50	S	E	5.05	5.9	5.1 5.4
April ..	30.035	5.9	16.7 3/25	57	S	E	4.05	5.0	4.3 7.8
May ..	30.083	5.8	17.9 17/26	48	SW	SE	3.09	4.1	4.3 8.3
June ..	30.091	5.7	19.0 14/28	58	SW	W & SW	2.45	2.9	4.4 9.2
July ..	30.090	5.6	15.0 2/23	52	SW	W & SW	2.69	2.8	3.8 12.4
August ..	30.105	5.8	14.8 4/35	56	SW	NE	3.51	3.8	3.1 13.1
September ..	30.067	5.9	16.1 1/48	57	SW	NE	4.51	5.8	3.3 13.0
October ..	30.019	6.3	15.7 1/41	62	S	NE	5.81	7.1	4.2 8.5
November ..	29.958	6.7	15.5 10/28	62	SE & N	NE	6.32	9.5	4.9 5.9
December ..	29.890	7.0	19.5 15/26	79	SE	NE	7.02	10.6	5.3 3.8
Year { Totals ..	—	—	—	—	—	—	56.73	73.8	— 93.3
Year { Averages ..	30.007	6.3	—	—	SW	NE	—	—	— 4.5
Year { Extremes ..	—	—	21.0 5/2/31	79	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940).

TEMPERATURE AND SUNSHINE.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).			Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	
							64		
No. of years over which observation extends.	30(a)	30(a)	30(a)	64	64	64	50(b)		30(a)
January ..	85.5	69.1	77.3	109.8 26/40	58.8 4/93	51.0	169.0 2/37	49.9 4/93	7.6
February ..	84.6	68.7	76.6	105.7 21/25	58.5 23/31	47.2	165.2 6/10	49.1 22/31	7.4
March ..	82.3	66.2	74.3	99.4 5/19	52.4 29/13	47.0	162.5 6/39	45.4 29/13	7.0
April ..	79.1	61.5	70.3	95.2 (c)	44.4 25/25	50.8	153.8 11/16	36.7 24/25	7.1
May ..	73.7	55.6	64.7	90.3 21/23	41.3 24/99	49.0	147.0 1/10	29.8 8/97	6.6
June ..	69.4	51.5	60.5	88.9 19/18	36.3 29/08	52.6	136.0 3/18	25.4 23/88	6.3
July ..	68.6	49.4	59.0	84.3 23/46	36.1 (d)	48.2	146.1 20/15	23.9 11/90	6.8
August ..	71.1	50.0	60.6	91.0 14/46	37.4 6/87	53.6	141.9 20/17	27.1 9/99	7.9
September ..	75.5	54.8	65.1	100.9 22/43	40.7 1/96	60.2	155.5 26/03	30.4 1/89	8.2
October ..	79.2	60.3	69.8	101.4 18/93	43.3 3/99	58.1	157.4 31/18	34.9 8/89	8.4
November ..	82.3	64.6	73.4	106.1 18/13	48.5 2/05	57.6	162.3 7/89	38.8 1/05	8.2
December ..	84.5	67.5	76.0	105.9 26/93	56.4 13/12	49.5	165.9 28/42	49.1 3/94	8.2
Year { Averages ..	78.0	59.9	69.0	—	—	—	—	—	7.5
Year { Extremes ..	—	—	—	109.8 26/1/40	36.1 (d)	73.7	169.0 2/1/37	23.9 11/7/90	—

(a) Standard 30 years' normal (1911-1940).

(b) From 1887 to March, 1947, excluding 1927 to 1936.

(c) 9/1896 and 5/1993.

(d) 12/7/1894 and 2/7/1896.

HUMIDITY, RAINFALL AND FOG.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).					Fog. Mean No. of Days of Fog.
		Mean.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.	
No. of years over which observation extends.	30(a)	30(a)	64	64	30(a)	30(a)	99		99	30(a)
January ..	0.636	66	79	53	5.72	12	27.72 1895	0.32 1919	18.31 21/87	0.6
February ..	0.644	69	82	55	5.47	12	40.39 1893	0.58 1849	10.61 6/31	0.9
March ..	0.606	72	85	56	4.97	14	34.04 1870	Nil 1849	11.18 14/08	1.6
April ..	0.512	71	80	56	3.68	11	35.28 1867	0.04 1944	5.46 5/33	4.0
May ..	0.420	71	85	59	2.35	9	13.85 1876	Nil 1846	5.62 9/79	5.4
June ..	0.357	73	84	54	2.75	8	14.03 1873	Nil 1847	6.41 15/48	4.9
July ..	0.331	71	88	53	1.88	8	8.60 1950	Nil 1841	4.89 1/4	4.9
August ..	0.338	67	80	55	1.07	7	14.67 1879	Nil (d)	4.89 12/87	2.8
September ..	0.396	62	76	47	1.69	7	5.43 1886	0.10 1907	2.46 2/94	2.8
October ..	0.459	59	72	48	2.27	8	11.41 1949	0.14 1900	5.34 25/49	1.6
November ..	0.533	61	72	45	4.00	10	12.40 1917	Nil 1842	4.46 16/86	0.7
December ..	0.589	62	70	51	4.24	11	17.36 1942	0.35 1865	6.60 28/71	0.4
Year { Totals ..	—	—	—	—	—	—	—	—	—	—
Year { Averages ..	0.485	67	—	—	40.09	117	—	—	—	33.3
Year { Extremes ..	—	—	85	45	—	—	40.39 2/1893	Nil (e)	18.31 21/1/87	—

(a) Standard 30 years' normal (1911-1940).

(b) Records incomplete for various years between 1846 and 1859.

(c) 15/1876 and 16/1889.

(d) 1862, 1869, 1880.

(e) Various months in various years.

CLIMATOLOGICAL DATA : SYDNEY, NEW SOUTH WALES.

LAT. 33° 52' S., LONG. 151° 12' E. HEIGHT ABOVE M.S.L. 138 FT.

BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS AND CLEAR DAYS.

Month.	Bar. corrected to 32° F. m. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 58 feet.)					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m. (a)	No. of Clear Days
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.					
					9 a.m.	3 p.m.				
No. of years of observations.	30(b)	26(c)	37 (d)	31 (e)	26(c)	26(c)	26(c)	30(f)	30(b)	30(b)
January ..	29.875	8.9	24.9 2/22	74	S	BNE	5.71	4.8	5.7	4.8
February ..	29.942	8.1	20.1 14/18	61	NE	BNE	4.68	3.3	5.5	5.4
March ..	30.009	7.5	20.7 10/44	58	W	BNE	4.05	2.8	5.3	5.8
April ..	30.063	7.0	23.4 19/27	72	W	NE	2.91	2.4	5.0	7.0
May ..	30.098	6.8	19.6 2/26	63	W	S	2.17	1.6	4.9	7.4
June ..	30.078	7.1	24.5 17/14	70	W	W	1.61	1.5	4.8	8.3
July ..	30.070	7.2	26.6 6/31	68	W	W	1.69	1.1	4.5	10.1
August ..	30.060	7.4	24.0 3/21	68	W	NE	2.30	2.1	3.9	11.1
September ..	30.018	8.0	22.3 19/17	70	W	NE	3.00	3.0	4.2	10.0
October ..	29.976	8.2	21.1 18/44	95	W	BNE	4.17	3.9	4.9	7.4
November ..	29.935	8.5	22.6 14/30	71	W & E	BNE	4.97	4.5	5.5	5.7
December ..	29.881	8.9	24.9 10/20	75	S	BNE	5.64	5.4	5.8	4.8
Year { Totals ..	30.000	7.8	26.6 6/7/31	95	W	NE	42.90	36.4	5.0	87.8
Year { Averages ..	—	—	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	—	—	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940). (c) 1915-1940. (d) 1914-1950. (e) 1917-1950. (f) 1921-1950.

TEMPERATURE AND SUNSHINE.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.	
No. of years over which observation extends.	30(a)	30(a)	30(a)	92	92	92	84	92	30(b)
January ..	78.6	65.1	71.8	113.6 14/39	51.1 18/49	62.5	164.3 26/15	43.7 6/25	7.5
February ..	78.7	65.5	72.1	107.8 8/25	49.3 28/63	58.5	168.3 14/39	42.8 22/33	7.0
March ..	76.0	62.9	69.8	102.6 3/59	48.8 14/86	53.8	158.3 10/26	39.9 17/13	6.4
April ..	72.0	57.7	64.9	91.4 3/36	44.6 27/64	46.8	144.1 10/77	33.3 24/09	6.1
May ..	67.0	52.4	59.7	86.0 1/19	40.2 22/59	45.8	129.7 1/66	29.3 25/17	5.7
June ..	62.8	48.1	55.5	80.4 11/31	35.7 22/32	44.7	125.5 2/33	28.0 22/32	5.3
July ..	61.8	46.4	54.1	78.3 22/26	35.9 12/90	42.4	124.7 19/77	24.0 4/93	6.1
August ..	64.3	47.6	56.0	82.8 12/46	36.8 3/72	46.0	140.0 30/78	26.1 4/09	7.0
September ..	68.3	51.4	59.9	92.3 27/19	40.8 3/45	51.5	142.2 12/78	30.1 17/05	7.3
October ..	71.7	55.9	63.8	99.4 4/42	42.2 6/27	57.2	152.2 20/33	32.7 9/05	7.5
November ..	74.5	59.8	67.1	104.5 6/46	45.8 1/05	61.7	158.5 28/39	36.0 6/06	7.5
December ..	76.9	63.2	70.1	107.5 31/04	48.4 3/24	59.1	164.5 27/89	41.4 3/24	7.5
Year { Averages ..	71.1	56.3	63.7	113.6 14/1/39	35.7 22/6/32	77.9	168.3 14/2/39	24.0 4/7/93	6.8
Year { Extremes ..	—	—	—	—	—	—	—	—	—

(a) Standard 30 years' normal (1911-1940). (b) 1921-1950 (different exposure prior to 1921).

HUMIDITY, RAINFALL AND FOG.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.				Rainfall (inches).				Fog. Mean No. of Days of Fog.
		Mean.	Highest.	Lowest.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.	
No. of years over which observation extends.	30(a)	30(a)	75	75	30(a)	30(a)	92	92	92	30(b)
January ..	0.537	65	78	58	3.86	13	15.26 1911	0.25 1932	7.08 13/11	0.4
February ..	0.560	68	81	60	3.15	12	18.56 1873	0.12 1939	8.90 25/73	0.8
March ..	0.527	71	85	62	4.44	13	20.52 1942	0.42 1876	11.05 28/42	1.8
April ..	0.441	73	87	63	5.65	14	24.49 1861	0.06 1868	7.52 29/60	2.8
May ..	0.362	75	90	63	4.98	12	23.03 1919	0.18 1860	8.36 28/89	3.7
June ..	0.303	76	89	65	3.68	11	25.30 1950	0.79 1904	5.17 16/84	3.3
July ..	0.282	74	88	63	4.89	12	13.23 1950	0.10 1946	7.80 7/31	2.9
August ..	0.288	68	84	54	2.41	10	14.89 1899	0.04 1885	5.33 2/60	2.3
September ..	0.325	62	79	49	2.77	11	14.05 1879	0.08 1882	5.69 10/79	1.0
October ..	0.378	60	77	46	2.80	11	11.13 1916	0.21 1867	6.37 13/02	0.6
November ..	0.433	60	79	42	2.54	11	9.88 1865	0.07 1915	4.23 19/00	0.6
December ..	0.501	63	77	51	3.63	13	15.82 1920	0.23 1913	4.75 13/10	0.4
Year { Totals ..	—	—	—	—	44.80	143	—	—	—	20.6
Year { Averages ..	0.393	68	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	90	42	—	—	25.30 6/1950	0.04 8/1885	11.05 28/3/42	—

(a) Standard 30 years' normal (1911-1940). (b) 1921-1950.

CLIMATOLOGICAL DATA : MELBOURNE, VICTORIA.

LAT. 37° 49' S., LONG. 144° 58' E. HEIGHT ABOVE M.S.L. 114 FT.

BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS AND CLEAR DAYS.

Month.	Bar. corrected to 32° F. Mm. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 93 feet.)					Mean Amount of Evaporation (Inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m. (a)	No. of Clear Days.	
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.						
					9 a.m.	3 p.m.					
No. of years of observations.	30(b)	11(c)	38	27	30(b)	30(b)	30(b)	30(b)	30(b)		
January	29.897	8.8	21.1	27/41	66	S & SW	S	6.55	1.8	4.9	6.8
February	29.950	8.3	19.0	13/47	66	N & S	S	5.10	2.3	4.8	6.4
March	30.025	7.8	17.2	19/50	66	N	S	4.26	1.8	5.3	5.5
April	30.092	7.2	19.9	16/43	67	N	S	2.53	1.2	5.0	4.6
May	30.113	7.4	20.0	4/44	72	N	S	1.57	0.5	6.1	3.4
June	30.097	7.4	22.8	16/47	60	N	S	1.18	0.4	6.5	2.7
July	30.079	8.6	20.9	9/44	68	N	S	1.16	0.3	6.3	2.9
August	30.048	8.2	21.3	20/42	64	N	S	1.54	0.9	6.0	3.1
September	30.001	8.5	18.3	6/48	69	N & W	N & S	2.41	1.3	5.9	3.3
October	29.968	8.3	18.0	27/50	69	N	S	3.54	1.6	6.1	3.6
November	29.951	8.4	19.4	4/50	65	S & SW	S	4.62	2.3	6.0	3.6
December	29.896	8.6	18.9	1/34	61	S & SW	S	5.85	1.9	5.6	4.5
Year { Totals	—	—	—	—	—	—	—	40.31	16.5	—	50.6
Year { Averages	30.010	8.1	—	—	—	N	S	—	—	5.8	—
Year { Extremes	—	—	22.8	16/6/47	72	—	—	—	—	—	—

(a) Scale 0-10.

(b) Standard 30 years' normal (1911-1940).

(c) Early records not comparable.

TEMPERATURE AND SUNSHINE.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.				
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.					
										30(a)	30(a)	30(a)	95
No. of years over which observation extends.	30(a)	30(a)	30(a)	95	95	95	86(b)	91	35(c)				
January	77.7	56.9	67.3	114.1	13/39	42.0	28/85	72.1	178.5	14/62	30.2	28/85	7.8
February	78.6	58.0	68.3	109.5	7/01	40.2	24/24	69.3	167.5	15/70	30.9	6/91	7.4
March	74.9	55.2	65.1	107.0	11/40	37.1	17/84	69.9	164.5	1/68	28.9	(d)	6.5
April	67.9	50.8	59.3	94.8	5/38	34.8	24/88	60.0	152.0	8/61	25.0	23/97	5.0
May	62.0	46.9	54.5	83.7	7/05	29.9	29/16	53.8	142.6	2/59	21.1	26/16	4.1
June	56.8	43.8	50.3	72.2	1/07	28.0	11/66	44.2	129.0	11/61	19.9	30/29	3.4
July	56.2	42.6	49.4	69.3	22/26	27.0	21/69	42.3	125.8	27/80	20.5	12/03	3.7
August	58.7	43.7	51.2	77.0	20/85	28.3	11/63	48.7	137.4	29/69	21.3	14/02	4.6
September	63.3	46.0	54.7	88.6	28/28	31.0	3/40	57.6	142.1	20/67	22.6	8/16	5.5
October	67.9	48.7	58.3	98.4	24/14	32.1	3/71	66.3	154.3	28/68	24.8	22/18	5.8
November	71.3	51.8	61.5	105.7	27/94	36.5	2/96	69.2	159.6	29/65	24.6	2/96	6.2
December	75.4	55.3	65.3	110.7	15/76	40.0	4/70	70.7	170.3	20/69	33.2	1/04	7.0
Year { Averages	67.6	50.0	58.8	—	—	—	—	—	—	—	—	—	5.6
Year { Extremes	—	—	—	114.1	13/1/39	27.0	21/7/60	87.1	178.5	14/1/62	19.9	30/6/29	—

(a) Standard 30 years' normal (1911-1940).

(b) Records discontinued, 1946.

(c) 1916-1930.

(d) 17/1884 and 20/1897.

HUMIDITY, RAINFALL AND FOG.

Month.	Vapour Pressure (Inches)	Rel. Hum. (%) at 9 a.m.				Rainfall (Inches).					Fog. Mean No. of Days of Fog.		
		Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.				
										30(a)		30(a)	43
No. of years over which observation extends.	30(a)	30(a)	43	43	30(a)	30(a)	95	95	95	30(a)			
January	0.382	58	65	50	1.88	9	6.66	1941	0.01	1932	2.97	9/97	0.1
February	0.417	62	70	48	2.00	8	7.72	1939	0.03	1870	3.44	26/46	0.3
March	0.385	64	76	50	2.22	9	7.50	1911	0.14	1934	3.55	5/19	1.1
April	0.351	72	82	66	2.30	13	6.71	1901	Nil	1923	2.28	22/01	2.3
May	0.311	79	86	70	1.94	14	5.60	1942	0.14	1934	1.85	7/91	6.8
June	0.276	83	92	75	2.06	16	4.51	1859	0.73	1877	1.74	21/04	6.5
July	0.264	82	86	75	1.93	17	7.02	1891	0.57	1902	2.71	12/91	6.5
August	0.271	76	82	65	2.02	17	4.35	1939	0.48	1903	1.94	26/24	3.7
September	0.288	68	76	60	2.20	15	7.93	1916	0.52	1907	2.62	12/80	1.3
October	0.307	62	67	52	2.63	14	7.61	1869	0.29	1914	3.00	17/69	0.3
November	0.336	60	69	52	2.33	13	6.71	1916	0.25	1895	2.57	16/76	0.3
December	0.373	59	69	48	2.38	11	7.18	1863	0.11	1904	3.20	1/34	0.2
Year { Totals	—	—	—	—	25.89	156	—	—	—	—	—	—	29.4
Year { Averages	0.323	69	—	—	—	—	—	—	—	—	—	—	—
Year { Extremes	—	—	92	48	—	—	7.93	9/1916	Nil	4/1923	3.55	5/3/19	—

(a) Standard 30 years' normal (1911-1940).

CLIMATOLOGICAL DATA : HOBART, TASMANIA.

LAT. 42° 53' S., LONG. 147° 30' E. HEIGHT ABOVE M.S.L. 177 FT.
 BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS AND CLEAR DAYS.

Month.	Bar. corrected to 32° F. Mm. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind.				Mean Amount of Evaporation (Inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m.(g)	No. of Clear Days.	
		(Height of Anemometer 40 feet.)								
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.					
			9 a.m.	3 p.m.						
No. of years of observations.	30(b)	30(b)	60	60	30(b)	30(b)	30(b)	30(b)	30(b)	
January ..	29.819	8.0	20.8 30/16	76	NNW	SSE	4.84	0.9	6.4	1.9
February ..	29.913	7.2	25.2 4/27	63	NNW	SSE	3.71	1.0	6.2	2.3
March ..	29.961	6.8	21.4 13/38	68	NW	SSE	3.10	1.2	6.1	2.4
April ..	29.997	6.7	22.2 27/26	74	NW	W	1.98	0.7	6.5	1.7
May ..	30.009	6.3	20.2 20/36	70	NNW	NW	1.37	0.4	6.1	2.4
June ..	29.986	6.2	23.7 27/20	71	NW	NW	0.91	0.4	6.2	2.4
July ..	29.958	6.5	20.8 19/35	78	NNW	NNW	0.94	0.3	6.1	2.0
August ..	29.906	6.8	25.5 19/26	87	NNW	NW	1.28	0.4	6.1	2.1
September ..	29.860	7.9	21.5 26/15	84	NNW	NW	1.97	0.7	6.3	1.5
October ..	29.833	8.2	19.2 8/12	74	NNW	SW	3.05	0.6	6.6	1.0
November ..	29.831	7.9	21.2 18/15	73	NNW	S	3.77	0.7	6.4	1.3
December ..	29.816	7.6	23.4 1/34	70	NNW	SSE	4.37	0.5	6.8	1.1
Year { Totals ..	—	—	—	—	—	—	31.29	7.8	—	22.1
{ Averages	29.907	7.2	—	—	NNW	W	—	—	6.3	—
{ Extremes	—	—	25.5 19/8/26	87	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940).

TEMPERATURE AND SUNSHINE.

Month.	Mean Temperature (°Fahr.)			Extreme Shade Temperature (°Fahr.).			Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	
	30(a)	30(a)	30(a)	67(b)	67(b)	67(b)	57(c)	67(b)	
No. of years over which observation extends.	30(a)	30(a)	30(a)	67(b)	67(b)	67(b)	57(c)	67(b)	30
January ..	69.8	52.4	61.0	105.0 1/00	40.1 (d)	64.9	160.0 (e)	30.6 19/97	7.7
February ..	70.6	53.7	62.2	104.4 12/99	39.0 20/87	65.4	165.0 24/98	28.3 —/87	7.1
March ..	67.5	51.3	59.4	99.1 13/40	35.2 31/26	63.9	150.9 26/44	27.5 30/02	6.4
April ..	62.2	48.0	55.1	87.1 1/41	33.3 24/88	53.8	142.0 18/93	25.0 —/86	5.0
May ..	57.8	44.6	51.2	77.8 5/21	29.2 20/02	48.6	128.0 (f)	20.0 19/02	4.4
June ..	52.8	41.2	47.0	69.2 1/07	29.2 28/44	40.0	122.0 12/94	21.0 6/87	4.0
July ..	52.7	40.6	46.6	66.1 14/34	27.7 11/95	38.4	121.0 12/93	18.7 16/86	4.4
August ..	55.4	41.7	48.7	71.6 28/14	30.5 (g)	41.1	129.0 —/87	20.1 7/09	5.1
September ..	59.0	43.7	51.4	81.7 23/26	31.0 16/97	50.7	138.0 23/93	18.3 16/26	5.9
October ..	62.5	46.1	54.3	92.0 24/14	32.0 12/89	60.0	156.0 9/93	23.8 (h)	6.1
November ..	65.0	48.2	56.6	98.3 26/37	35.0 16/41	63.3	154.0 19/92	26.0 1/08	7.2
December ..	67.9	51.3	59.6	105.2 30/97	38.0 3/06	67.2	161.5 10/39	27.2 —/86	7.3
Year { Averages	61.9	46.9	54.4	—	—	—	—	—	5.9
{ Extremes	—	—	—	105.2 30/12/97	27.7 11/7/95	77.5	165.0 24/2/98	18.3 16/9/26	—

(a) Standard 30 years' normal (1911-1940). (b) Records 1855-1882 not comparable. (c) Period 1934-1938 not comparable; records discontinued, 1946. (d) 9/37 and 11/37. (e) 5/86 and 13/05. (f) —/89 and —/93. (g) 4/97 and 7/09. (h) 1/86 and —/99.

HUMIDITY, RAINFALL AND FOG.

Month.	Vapour Pressure (Inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (Inches).				Fog.	
		Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.		Greatest in One Day.
No. of years over which observation extends.	30(a)	31	64	64	30(a)	30(a)	67(b)	67(b)	67(b)	29
January ..	0.309	57	72	46	1.82	13	5.91 1803	0.17 1915	2.96 30/16	0.0
February ..	0.342	61	77	48	1.68	10	4.96 1935	0.11 1914	2.18 5/38	0.0
March ..	0.323	65	77	52	2.13	13	10.05 1946	0.29 1943	3.47 17/46	0.3
April ..	0.290	69	84	58	2.31	14	8.50 1935	0.07 1904	5.02 20/09	0.2
May ..	0.263	78	89	65	1.71	14	6.37 1905	0.14 1913	1.75 2/93	0.9
June ..	0.233	78	91	68	2.25	16	8.15 1889	0.28 1886	4.11 13/89	0.7
July ..	0.227	78	94	72	2.14	17	6.02 1922	0.17 1950	2.51 18/22	0.9
August ..	0.232	72	92	60	1.82	18	6.32 1946	0.30 1802	2.28 14/90	0.4
September ..	0.240	64	85	58	1.90	17	4.47 1928	0.40 1801	1.57 24/85	0.1
October ..	0.258	60	73	51	2.52	18	7.60 1947	0.39 1914	2.58 4/06	0.0
November ..	0.274	57	72	50	2.23	16	7.39 1885	0.33 1921	3.70 30/85	0.0
December ..	0.306	58	67	45	2.52	14	7.72 1916	0.17 1931	3.33 5/41	0.0
Year { Totals ..	—	—	—	—	25.03	180	—	—	—	3.5
{ Averages	0.271	66	—	—	—	—	—	—	—	—
{ Extremes	—	—	94	45	—	—	10.05 3/1946	0.07 4/1904	5.02 20/4/09	—

(a) Standard 30 years' normal (1911-1940). (b) Records prior to 1883 not comparable.

§ 3. Standard Times in Australia.

Prior to 1895 the official time adopted in the several colonies was for most purposes the mean solar time of the capital city of each.

In November, 1892, an intercolonial conference of surveyors was held in Melbourne to consider, among other things, the advantages of introducing the system of standard time. In this system it was proposed to make the initial meridian that of Greenwich and to change local standard time by whole hours according to the longitude east or west of that of Greenwich. Thus for every difference of 15° in longitude a change of one hour would be required. The minutes and seconds would then be identical everywhere.

To give effect to this proposal it was suggested that Australia should be divided into three zones, the standard times for which should be respectively the mean solar times of the meridians of 120° , 135° and 150° E. longitude, thus giving standard times 8, 9 and 10 hours respectively ahead of Greenwich time. It was proposed that the 120° zone should comprise Western Australia, that the 135° zone should comprise South Australia and the Northern Territory, and that the 150° zone should comprise Queensland, New South Wales, Victoria and Tasmania.

The matter was also considered by several intercolonial postal conferences, and eventually in 1894 and 1895 legislation was enacted by each of the colonies in accord with the recommendations of the Surveyors' Conference of 1892.

In 1898 the South Australian legislature amended its earlier provision, and adopted the mean solar time of the meridian $142^\circ 30'$ E. longitude as the standard time for that colony, thus reducing the difference between the standard time of Adelaide and that of the capitals of the eastern colonies from an hour to half-an-hour. Particulars concerning these enactments are as follows :—

STANDARD TIMES IN AUSTRALIA.

State.	Date when Act came into Operation.	Meridian Selected.	Time Ahead of Greenwich.
			Hours.
New South Wales ..	1st February, 1895 ..	150° E.	10
Victoria ..	1st February, 1895 ..	150° E.	10
Queensland ..	1st January, 1895 ..	150° E.	10
South Australia ..	1st February, 1895 ..	135° E.	9
South Australia ..	1st May, 1899 ..	$142^\circ 30'$ E.	$9\frac{1}{2}$
Western Australia ..	1st December, 1895 ..	120° E.	8
Tasmania ..	1st September, 1895 ..	150° E.	10

The standard time in the Australian Capital Territory is the same as in New South Wales, and in the Northern Territory the same as in South Australia.

Consequent upon the opening of the Trans-Australian Railway an arrangement was made by which the change of time between South Australia and Western Australia (namely, $1\frac{1}{2}$ hours) is divided into two changes of 45 minutes each. Going east from Kalgoorlie the first change is made at Rawlinna, 235.18 miles out, where the time is put forward by 45 minutes. The second change of the same amount is made at Tarcoola, 794.05 miles out. Thenceforward South Australian standard time is kept. The Commonwealth Observatory at Mount Stromlo, Canberra, and the State Observatories at Sydney and Perth derive time by astronomical observations.

Time signals are originated by these Observatories and by the Postmaster-General's Research Laboratory, Melbourne. The latter participates with the Commonwealth Observatory in the Commonwealth Time Service.