



INFORMATION PAPER

SUICIDES 2001

TABLE OF CONTENTS

INTRODUCTION	3
SUICIDE BY YEAR OF REGISTRATION	4
SUICIDE BY AGE AND SEX, Number of Deaths	9
SUICIDE BY AGE AND SEX, Age-Specific Death Rates	10
SUICIDE BY METHOD, Number of Deaths	11
SUICIDE BY METHOD, Proportion of Total Suicides	12
SUICIDE AND ALL CAUSES, Number of Deaths and Years of Potential Life Lost	13
SUICIDE BY STATE OR TERRITORY AND SEX, Number of Deaths	14
SUICIDE BY STATE OR TERRITORY AND SEX, Standardised Death Rates	15
SUICIDE BY YEAR OF OCCURRENCE	16
SUICIDE BY AGE AND SEX, Number of Deaths	16
SUICIDE BY AGE AND SEX, Age-Specific Death Rates	17
SUICIDES IN THE 15-19 & 20-24 YEAR AGE GROUPS	18
SUICIDE BY AGE AND SEX, Number and Rates - Year of registration	18
SUICIDE BY AGE AND SEX, Number and Rates - Year of occurrence	19
APPENDIX	20

INTRODUCTION

Suicide continues to be a major public health issue. Although death by suicide is a relatively uncommon event (in 2001, only 1.9% of all deaths registered were attributed to suicide), the human and economic costs are significant. Apart from the loss of life, there are health care costs associated with attempted suicide and the circumstances surrounding death can be particularly difficult for family and friends to deal with.

Suicide can be defined as the deliberate taking of one's life. To be classified as a suicide, a death must be recognised as due to other than natural causes. It must also be established by coronial inquiry that the death results from a deliberate act of the deceased with the intent of ending his or her own life. Possible suicides where the coroner concludes an 'open' finding and not explicitly suicide are excluded. See explanatory notes in Suicides 1921-1998 (ABS Cat. No. 3309.0) for further explanation of deaths of undetermined intent.

This paper contains summary statistics where the underlying cause of death was determined as suicide. Initially data are presented by year of registration. A later section presents data by year of occurrence (see page 16). It is common international practice when publishing data on a year of occurrence basis to include deaths registered in the particular occurrence year and the year immediately following. This practice has been followed in this paper.

In this paper, suicide deaths data for registration years 1991-1996 were coded to ICD-9 while data for registration years 1997-2001 were coded to ICD-10. For suicide deaths, ICD-9 and ICD-10 classifications are comparable.

Other ABS publications or articles which contain information on suicides are as follows:

Publications:

Suicides 1921-1998 (ABS Cat. No. 3309.0)

Causes of death, Australia (ABS Cat. No. 3303.0)- annual

Deaths, Australia 3302.0 (ABS Cat. No. 3302.0) - annual

A Mortality Atlas of Australia (ABS Cat. No. 3318.0)

Trends in mortality by causes of death in Australia, the states and territories during 1971-92, and in statistical divisions and sub-divisions during 1991-92 (ABS Cat. No. 3313.0)

Articles:


Suicide (Australian Social Trends 2000, ABS Cat. No. 4102.0)

Youth suicide (Australian Social Trends 1994, ABS Cat. No. 4102.0)

SUICIDE BY YEAR OF REGISTRATION

For 2001, (the latest year available) 2454 suicides were registered, compared with 2363 for the year 2000. The age standardised rate for the year 2001 (12.5 deaths per 100,000 persons) was 15% lower than it was in 1997 (14.6 deaths per 100,000) when the highest ever recorded number of 2720 suicides were registered; and 2.2% higher than the 2000 age standardised rate of 12.2 per 100,000 persons.

AGE-STANDARDISED DEATH RATES FOR SUICIDE



AGE AND SEX

Over the period 1991 to 2001 (registration years) male suicides outnumbered female suicides by approximately four to one. This ratio was fairly consistent over all age groups. While publicity has focussed on youth suicide in recent years, the age group with the highest age specific death rate in 2001 was the 25-34 year age group with 20.6 per 100,000, followed by the 35-44 year age group with a rate of 18.9 per 100,000. Numerically these two age groups accounted for 47% of total suicide deaths in 2001.

Major figures for deaths in each age group by registration year include:

15 - 24 year age group

- In 2001, suicide accounted for 339 deaths with a death rate of 12.7 per 100,000 persons.
- In 2001, suicide accounted for 25% of total male deaths and 16% of total female deaths in this age group.
- Over the period 1991-2001, the lowest number of suicides registered in this age group was 338 in 2000 (age-specific rate of 12.8), while the highest number of suicides was 509 deaths registered in 1997 (rate of 19.3).

25 - 34 year age group

- In 2001, suicide accounted for 594 deaths with a death rate of 20.6 per 100,000 persons.
- In 2001, suicide accounted for 30% of total male deaths and 18% of total female deaths in this age group
- Over the period 1991-2001 suicides registered varied from a low of 484 deaths (rate of 17.1) in 1993 to a high of 700 deaths (rate of 24.2) in 1998.

35 - 44 year age group

- In 2001, suicide accounted for 565 deaths with a death rate of 18.9 per 100,000 persons.
- In 2001, suicide accounted for 20% of total male deaths and 9% of total female deaths.
- Over the period 1991-2001 suicides registered varied from a low of 366 deaths (rate of 13.7) in 1993 to a high of 614 deaths (rate of 21.1) in 1998.

45 - 54 year age group

- In 2001, suicide accounted for 417 deaths with a death rate of 15.6 per 100,000 persons.
- In 2001, suicide accounted for 8% of total male deaths and 4% of total female deaths.
- Over the period 1991 - 2001 suicides registered varied from a low of 318 deaths (rate of 15.4) in 1993 to a high of 417 deaths (rate of 15.6) in 2001.

55 - 64 year age group

- In 2001, suicide accounted for 223 deaths with a death rate of 12.1 per 100,000 persons.
- In 2001, suicide accounted for 2% of total male deaths and 1% of total female deaths.
- Over the period 1991 - 2001 suicides registered varied from a low of 197 deaths (rate of 11.1) in 2000 to a high of 233 deaths (rate of 14.7) in 1997.

65 - 74 year age group


- In 2001, suicide accounted for 160 deaths with a death rate of 12.1 per 100,000 persons.
- In 2001, suicide accounted for less than 1% of total male deaths and of total female deaths.

75+ year age group

- In 2001, suicide accounted for 145 deaths with a death rate of 13.0 per 100,000 persons.
- In 2001, suicide accounted for less than 1% of total male deaths and of total female deaths.

Detailed information on suicide by sex and age groups (numbers and rates) for 1991 - 2001 is included in Tables 1 and 2

AGE-SPECIFIC DEATH RATES FOR SUICIDE




METHOD OF SUICIDE

Over registration years 1991 - 2001, there have been noticeable changes in the various methods employed for suicide. These include:

- A decrease in the use of firearms. Over this period, the proportion of male suicides using this method decreased from 26% of total male suicides in 1991 to 13% in 2001, while the corresponding female proportion decreased from 4.5% in 1991 to 3.9% in 2001.
- An increase in hanging, strangulation and suffocation. Over this period, the proportion of male suicides using this method increased from 26% of total male suicides in 1991 to 44% in 2001, while the corresponding female proportion increased from 21% in 1991 to 38% in 2001.
- A decrease in poisoning by drugs and medicaments. Over this period, the proportion of female suicides using this method decreased from 35% of total female suicides in 1991 to 26% in 2001, while the corresponding male proportion decreased from 11% in 1991 to 7.8% in 2001.
- The proportion of suicides where the method was “poisoning by other” (primarily motor vehicle exhaust) has remained relatively stable over this period.

Detailed information on method of suicide (numbers and rates) for 1991 to 2001 is included in Tables 3 and 4.

SUICIDE BY METHOD, 1991 AND 2001



(a) Includes other solids, liquids, gases and vapours (including motor vehicle exhaust)

(b) Includes strangulation and suffocation.

(c) Includes drowning, cutting and piercing instruments, jumping from high places and other and unspecified means.

YEARS OF POTENTIAL LIFE LOST (YPLL)

Years of Potential Life Lost (YPLL) is a method used to measure the extent of premature mortality associated with various causes of mortality. This measurement assumes that premature mortality is any death between ages 1 and 75 years for both males and females. In registration year 2001:

- Male suicide accounted for 5.5% of total male deaths 1 to 75 years of age but was responsible for 13% of the total Years of Potential Life Lost.
- Female suicide accounted for 2.5% of total female deaths 1 to 75 years of age but was responsible for 6% of the total Years of Potential Life Lost.
- Suicide as a proportion of total YPLL has risen from 8.7% to 10.6% over the period 1991-2001

More detailed information on Years of Potential Life Lost for suicide is included in Table 5.

STATE AND TERRITORY

Over the eleven year period 1991 to 2001, there was some variation in the standardised suicide death rate (for year of registration) across States and Territories, including:

- The Queensland suicide death rate has been higher than the national rate for each of the eleven years.
- Suicide death rates for Northern Territory and Tasmania have been higher than the national rate for ten and eight years respectively.
- Victoria's suicide death rate has been lower than the national rate for each of the eleven years.
- Suicide death rates for New South Wales and Australian Capital Territory have been lower than the national rate for seven and nine years respectively.

Detailed information on suicide rates by sex for States and Territories for registration years 1991 to 2001 is included in Tables 6 and 7.

1. SUICIDE BY AGE AND SEX, Number of Deaths

Year of registration	15-24	25-34	35-44	45-54	55-64	65-74	75+	All Ages(a)
MALES								
1991	376	423	400	251	156	125	110	1,847
1992	378	431	331	260	170	154	91	1,820
1993	345	406	286	247	170	135	94	1,687
1994	374	414	354	270	173	141	100	1,830
1995	350	475	382	270	177	117	96	1,873
1996	351	465	413	266	181	136	112	1,931
1997	416	540	431	294	177	146	131	2,143
1998	364	591	479	314	163	136	97	2,150
1999	309	507	427	312	176	140	121	2,002
2000	263	475	444	291	146	120	113	1,860
2001	276	486	447	311	175	127	107	1,935
FEMALES								
1991	86	104	96	71	73	42	39	513
1992	77	100	95	62	52	45	40	474
1993	55	78	80	71	37	39	33	394
1994	57	85	99	68	47	32	37	428
1995	84	99	104	89	50	37	32	495
1996	56	96	110	88	40	38	27	462
1997	93	115	122	96	56	47	41	577
1998	82	109	135	72	53	41	39	533
1999	71	116	104	97	42	29	23	490
2000	75	106	132	67	51	41	30	503
2001	63	108	118	106	48	33	38	519
PERSONS								
1991	462	527	496	322	229	167	149	2,360
1992	455	531	426	322	222	199	131	2,294
1993	400	484	366	318	207	174	127	2,081
1994	431	499	453	338	220	173	137	2,258
1995	434	574	486	359	227	154	128	2,368
1996	407	561	523	354	221	174	139	2,393
1997	509	655	553	390	233	193	172	2,720
1998	446	700	614	386	216	177	136	2,683
1999	380	623	531	409	218	169	144	2,492
2000	338	581	576	358	197	161	143	2,363
2001	339	594	565	417	223	160	145	2,454

(a) Includes deaths aged under 15 and age not stated.

2. SUICIDE BY AGE AND SEX, Age-Specific Death Rates (a)

Year of registration	15-24	25-34	35-44	45-54	55-64	65-74	75+	All Ages (b)
MALES								
1991	26.7	29.9	30.3	26.1	21.3	22.8	38.2	21.7
1992	27.0	30.4	24.9	25.8	23.1	27.3	30.6	21.1
1993	24.7	28.7	21.4	23.5	22.9	23.3	30.6	19.3
1994	27.0	29.2	26.1	24.7	23.1	23.6	31.7	20.7
1995	25.4	33.4	27.8	23.9	23.3	19.3	29.2	20.9
1996	25.7	32.5	29.4	22.7	23.4	22.2	32.4	21.3
1997	30.9	37.5	30.2	24.3	22.2	23.5	36.1	23.5
1998	27.3	41.1	33.2	25.1	19.9	21.7	25.6	23.2
1999	23.2	35.3	29.2	24.4	20.6	22.2	30.5	21.3
2000	19.6	33.1	30.2	22.3	16.4	18.9	27.2	19.5
2001	20.3	33.9	30.2	23.3	18.9	19.9	24.4	20.1
FEMALES								
1991	6.3	7.4	7.4	7.8	10.0	6.6	8.1	5.9
1992	5.7	7.1	7.2	6.4	7.1	7.0	8.1	5.3
1993	4.1	5.5	6.0	7.1	5.0	5.9	6.5	4.4
1994	4.3	6.0	7.3	6.5	6.3	4.8	7.1	4.7
1995	6.4	7.0	7.5	8.1	6.6	5.5	5.9	5.4
1996	4.3	6.7	7.8	7.7	5.2	5.6	4.8	4.9
1997	7.2	8.0	8.5	8.1	7.1	6.9	7.0	6.1
1998	6.4	7.5	9.2	5.9	6.5	6.0	6.4	5.5
1999	5.5	8.0	7.0	7.7	5.0	4.3	3.6	5.1
2000	5.8	7.3	8.8	5.1	5.8	6.0	4.6	5.1
2001	4.8	7.4	7.8	7.9	5.3	4.8	5.6	5.1
PERSONS								
1991	16.7	18.7	18.9	17.2	15.7	14.1	19.4	13.7
1992	16.5	18.7	16.1	16.3	15.1	16.5	16.5	13.1
1993	14.6	17.1	13.7	15.4	14.0	14.0	15.5	11.7
1994	15.9	17.6	16.7	15.8	14.7	13.6	16.4	12.6
1995	16.1	20.2	17.6	16.1	15.0	12.0	14.7	13.0
1996	15.2	19.6	18.6	15.3	14.4	13.4	15.3	13.0
1997	19.3	22.7	19.3	16.3	14.7	14.8	18.1	14.6
1998	17.0	24.2	21.1	15.6	13.2	13.5	13.8	14.3
1999	14.5	21.6	18.1	16.1	12.9	12.9	14.0	13.0
2000	12.8	20.1	19.4	13.7	11.1	12.2	13.4	12.2
2001	12.7	20.6	18.9	15.6	12.1	12.1	13.0	12.5

(a) Deaths per 100,000 of the estimated mid-year population for each age and sex group.

(b) Standardised death rate per 100,000 persons. Standardised to the estimated mid-year 1991 Australian population.

3. SUICIDE BY METHOD, Number of Deaths

Year of registration	Poisoning by drugs	Poisoning by other (a)	Hanging (b)	Firearms and explosives	Other (c)	Total
MALES						
1991	208	448	479	487	225	1,847
1992	172	459	496	464	229	1,820
1993	135	395	507	418	232	1,687
1994	188	425	532	400	285	1,830
1995	194	455	585	366	273	1,873
1996	163	486	672	369	241	1,931
1997	160	555	812	309	307	2,143
1998	157	499	1,035	218	241	2,150
1999	158	492	868	257	227	2,002
2000	144	472	807	213	224	1,860
2001	151	427	855	242	260	1,935
FEMALES						
1991	180	87	108	23	115	513
1992	179	68	93	26	108	474
1993	127	67	88	17	95	394
1994	142	72	107	20	87	428
1995	154	101	114	23	103	495
1996	165	94	120	15	68	462
1997	149	116	175	21	116	577
1998	153	87	182	17	94	533
1999	120	103	160	13	94	490
2000	129	102	182	10	80	503
2001	134	86	195	20	84	519
PERSONS						
1991	388	535	587	510	340	2,360
1992	351	527	589	490	337	2,294
1993	262	462	595	435	327	2,081
1994	330	497	639	420	372	2,258
1995	348	556	699	389	376	2,368
1996	328	580	792	384	309	2,393
1997	309	671	987	330	423	2,720
1998	310	586	1,217	235	335	2,683
1999	278	595	1,028	270	321	2,492
2000	273	574	989	223	304	2,363
2001	285	513	1,050	262	344	2,454

(a) Includes other solids and liquids and gases and vapours (including motor vehicle exhaust).

(b) Includes strangulation and suffocation.

(c) Includes drowning, cutting and piercing instruments, jumping from high places and other and unspecified means.

4. SUICIDE BY METHOD, Proportion of Total Suicides

Year of registration	Poisoning by drugs	Poisoning by other (a)	Hanging (b)	Firearms and explosives	Other (c)	Total
MALES						
1991	11.3	24.3	25.9	26.4	12.2	100.0
1992	9.5	25.2	27.3	25.5	12.6	100.0
1993	8.0	23.4	30.1	24.8	13.8	100.0
1994	10.3	23.2	29.1	21.9	15.6	100.0
1995	10.4	24.3	31.2	19.5	14.6	100.0
1996	8.4	25.2	34.8	19.1	12.5	100.0
1997	7.5	25.9	37.9	14.4	14.3	100.0
1998	7.3	23.2	48.1	10.1	11.2	100.0
1999	7.9	24.6	43.4	12.8	11.3	100.0
2000	7.7	25.4	43.4	11.5	12.0	100.0
2001	7.8	22.1	44.2	12.5	13.4	100.0
FEMALES						
1991	35.1	17.0	21.1	4.5	22.4	100.0
1992	37.8	14.3	19.6	5.5	22.8	100.0
1993	32.2	17.0	22.3	4.3	24.1	100.0
1994	33.2	16.8	25.0	4.7	20.3	100.0
1995	31.1	20.4	23.0	4.6	20.8	100.0
1996	35.7	20.3	26.0	3.2	14.7	100.0
1997	25.8	20.1	30.3	3.6	20.1	100.0
1998	28.7	16.3	34.1	3.2	17.6	100.0
1999	24.5	21.0	32.7	2.7	19.2	100.0
2000	25.6	20.3	36.2	2.0	15.9	100.0
2001	25.8	16.6	37.6	3.9	16.2	100.0
PERSONS						
1991	16.4	22.7	24.9	21.6	14.4	100.0
1992	15.3	23.0	25.7	21.4	14.7	100.0
1993	12.6	22.2	28.6	20.9	15.7	100.0
1994	14.6	22.0	28.3	18.6	16.5	100.0
1995	14.7	23.5	29.5	16.4	15.9	100.0
1996	13.7	24.2	33.1	16.0	12.9	100.0
1997	11.4	24.7	36.3	12.1	15.6	100.0
1998	11.6	21.8	45.4	8.8	12.5	100.0
1999	11.2	23.9	41.3	10.8	12.9	100.0
2000	11.6	24.3	41.9	9.4	12.9	100.0
2001	11.6	20.9	42.8	10.7	14.0	100.0

(a) Includes other solids and liquids and gases and vapours (including motor vehicle exhaust).

(b) Includes strangulation and suffocation.

(c) Includes drowning, cutting and piercing instruments, jumping from high places and other and unspecified means.

5. SUICIDE AND ALL CAUSES, Number of Deaths and Years of Potential Life Lost (YPLL) (a)

Year of registration	Number of deaths (b)		Years of Potential Life Lost (c)		SUICIDE AS A PERCENTAGE OF TOTAL	
	<i>Suicide</i>	<i>All Causes</i>	<i>Suicide</i>	<i>All Causes</i>	<i>Deaths (b)</i>	<i>YPLL</i>
1991	2,228	59,784	81,371	937,178	3.7	8.7
1992	2,170	60,386	78,873	930,112	3.6	8.5
1993	1,966	58,928	70,912	907,161	3.3	7.8
1994	2,135	59,683	77,802	902,319	3.6	8.6
1995	2,252	58,994	82,974	902,900	3.8	9.2
1996	2,270	58,914	83,407	897,389	3.9	9.3
1997	2,565	58,472	96,653	899,640	4.4	10.7
1998	2,559	56,782	97,074	884,189	4.5	11.0
1999	2,361	55,632	87,637	864,679	4.2	10.1
2000	2,235	54,232	82,922	836,594	4.1	9.9
2001	2,324	53,351	85,324	801,266	4.4	10.6

(a) YPLL measures the extent of premature mortality which is assumed to be any death occurring between ages 1 and 75 inclusive.

(b) Deaths occurring between ages 1-75 years inclusive.

(c) YPLL age-standardised to June 1991 population - see Technical Note in ABS 2001 Causes of Death Australia 3303.0.

6. SUICIDE BY STATE OR TERRITORY(a) AND SEX, Number of Deaths

Year of registration	NSW	Vic	Qld	SA	WA	Tas	NT	ACT	Aust
MALES									
1991	593	472	345	172	170	53	16	26	1,847
1992	586	445	331	165	172	75	20	26	1,820
1993	567	403	305	132	173	67	17	23	1,687
1994	629	401	365	140	188	63	18	26	1,830
1995	593	439	396	161	179	51	22	31	1,873
1996	660	388	451	153	171	54	28	26	1,931
1997	712	528	434	161	206	40	33	29	2,143
1998	705	446	454	199	234	49	35	28	2,150
1999	689	440	396	155	192	68	27	35	2,002
2000	588	382	421	161	206	40	37	25	1,860
2001	613	412	406	168	212	51	39	34	1,935
FEMALES									
1991	174	135	78	59	42	14	3	8	513
1992	146	112	97	48	42	21	3	5	474
1993	136	94	62	34	43	16	5	4	394
1994	152	110	90	29	29	7	1	10	428
1995	172	127	99	39	39	15	1	3	495
1996	155	112	88	32	47	10	7	11	462
1997	222	141	101	35	49	11	5	13	577
1998	157	133	125	45	53	10	7	3	533
1999	180	112	84	45	44	10	5	10	490
2000	142	129	120	38	55	10	5	4	503
2001	172	129	93	39	57	13	4	12	519
PERSONS									
1991	767	607	423	231	212	67	19	34	2,360
1992	732	557	428	213	214	96	23	31	2,294
1993	703	497	367	166	216	83	22	27	2,081
1994	781	511	455	169	217	70	19	36	2,258
1995	765	566	495	200	218	66	23	34	2,368
1996	815	500	539	185	218	64	35	37	2,393
1997	934	669	535	196	255	51	38	42	2,720
1998	862	579	579	244	287	59	42	31	2,683
1999	869	552	480	200	236	78	32	45	2,492
2000	730	511	541	199	261	50	42	29	2,363
2001	785	541	499	207	269	64	43	46	2,454

(a) State or Territory of Usual Residence

7. SUICIDE BY STATE OR TERRITORY(a) AND SEX, Standardised Death Rates (b)

Year of registration	NSW	Vic	Qld	SA	WA	Tas	NT	ACT	Aust
MALES									
1991	20.4	21.6	23.9	24.0	21.0	23.7	17.8	19.0	21.7
1992	19.9	20.2	22.3	22.6	21.0	33.1	25.7	18.2	21.1
1993	19.1	18.2	20.1	18.1	20.7	29.4	26.6	14.4	19.3
1994	21.1	18.0	23.2	19.1	22.1	27.7	17.9	16.6	20.7
1995	19.5	19.5	24.3	22.3	20.7	22.4	22.5	22.6	20.9
1996	21.6	17.1	27.3	20.5	19.4	24.0	28.3	16.9	21.3
1997	23.0	23.2	26.0	21.9	23.2	17.4	36.5	18.7	23.5
1998	22.6	19.3	26.6	27.3	26.2	20.9	35.8	19.4	23.2
1999	21.5	18.9	22.7	20.9	21.1	30.0	24.5	22.7	21.3
2000	18.1	16.2	24.1	21.3	21.8	18.5	33.4	15.6	19.5
2001	18.4	17.3	22.8	22.7	22.7	22.8	39.3	21.2	20.1
FEMALES									
1991	5.8	6.0	5.3	8.0	5.2	6.0	2.8	5.5	5.9
1992	4.7	4.9	6.4	6.5	5.1	8.7	3.6	3.8	5.3
1993	4.4	4.1	4.0	4.5	5.2	6.6	5.6	2.6	4.4
1994	4.8	4.7	5.6	3.9	3.5	3.0	6.4	6.1	4.7
1995	5.5	5.5	6.0	5.1	4.5	6.3	4.9	2.0	5.4
1996	4.8	4.7	5.2	4.2	5.3	3.8	7.8	6.6	4.9
1997	6.8	5.9	5.9	4.5	5.5	4.4	5.1	8.5	6.1
1998	4.8	5.6	7.1	5.4	5.8	3.7	7.2	1.9	5.5
1999	5.5	4.6	4.8	5.8	4.7	4.5	5.0	5.4	5.1
2000	4.2	5.2	6.6	5.0	5.9	4.5	5.0	2.0	5.1
2001	4.9	5.1	5.0	4.8	5.7	5.1	4.0	7.0	5.1
PERSONS									
1991	13.0	13.6	14.5	15.8	13.1	14.6	10.9	11.8	13.7
1992	12.2	12.4	14.3	14.5	13.0	20.7	15.5	10.6	13.1
1993	11.6	11.0	11.8	11.1	13.0	18.0	16.4	8.4	11.7
1994	12.8	11.2	14.3	11.4	12.8	15.0	13.1	11.3	12.6
1995	12.4	12.3	15.1	13.6	12.6	14.2	14.7	11.7	13.0
1996	13.0	10.8	16.2	12.2	12.3	13.5	18.9	11.8	13.0
1997	14.8	14.3	15.8	13.1	14.2	10.7	21.5	13.5	14.6
1998	13.5	12.3	16.8	16.4	15.9	12.1	22.3	10.1	14.3
1999	13.3	11.5	13.7	13.3	12.8	16.9	15.2	13.7	13.0
2000	11.0	10.5	15.2	13.1	13.9	11.4	19.9	8.8	12.2
2001	11.5	11.0	13.8	13.7	14.2	13.6	22.6	14.0	12.5

(a) State or Territory of Usual Residence.

(b) Standardised death rate per 100,000 persons. Standardised to the estimated mid year 1991 Australian population.

SUICIDES BY YEAR OF OCCURRENCE

8. SUICIDE BY AGE AND SEX, Number of Deaths

Year of occurrence(a)	15-24	25-34	35-44	45-54	55-64	65-74	75+	All Ages(b)
MALES								
1991	354	427	379	252	142	131	114	1,806
1992	384	407	328	251	175	150	87	1,786
1993	332	392	282	241	164	129	95	1,641
1994	377	426	366	282	173	138	100	1,865
1995	342	475	383	268	171	120	91	1,854
1996	366	469	408	259	186	144	122	1,962
1997	407	529	425	292	171	132	123	2,086
1998	353	579	466	307	159	142	96	2,108
1999	303	511	431	310	180	135	119	1,999
2000	270	465	449	295	151	129	115	1,883
2001	na	na	na	na	na	na	na	na
FEMALES								
1991	84	91	103	66	63	33	39	482
1992	76	96	83	61	54	48	45	466
1993	50	76	87	74	37	38	34	398
1994	55	88	97	65	48	31	33	418
1995	85	100	111	97	46	37	29	505
1996	58	98	105	77	47	39	28	459
1997	86	109	121	98	49	47	40	557
1998	81	109	128	74	52	38	40	525
1999	70	117	107	94	46	28	21	489
2000	78	106	132	68	47	40	33	505
2001	na	na	na	na	na	na	na	na
PERSONS								
1991	438	518	482	318	205	164	153	2,288
1992	460	503	411	312	229	198	132	2,252
1993	382	468	369	315	201	167	129	2,039
1994	432	514	463	347	221	169	133	2,283
1995	427	575	494	365	217	157	120	2,359
1996	424	567	513	336	233	183	150	2,421
1997	493	638	546	390	220	179	163	2,643
1998	434	688	594	381	211	180	136	2,633
1999	373	628	538	404	226	163	140	2,488
2000	348	571	581	363	198	169	148	2,388
2001	na	na	na	na	na	na	na	na

(a) Data for each year of occurrence includes deaths registered in the particular occurrence year and the year immediately following.

(b) Includes deaths aged under 15 and age not stated.

na not available

9. SUICIDE BY AGE AND SEX, Age-Specific Death Rates (a)

Year of occurrence (c)	15-24	25-34	35-44	45-54	55-64	65-74	75+	All Ages (b)
MALES								
1991	25.2	30.1	28.7	26.2	19.3	23.9	39.6	21.3
1992	27.4	28.7	24.7	24.9	23.7	26.6	29.2	20.7
1993	23.8	27.7	21.1	22.9	22.1	22.2	31.0	18.8
1994	27.2	30.1	27.0	25.8	23.1	23.1	31.7	21.1
1995	24.9	33.4	27.8	23.7	22.5	19.8	27.7	20.7
1996	26.8	32.8	29.1	22.1	24.0	23.5	35.3	21.7
1997	30.3	36.8	29.8	24.2	21.5	21.3	33.9	22.8
1998	26.4	40.2	32.3	24.6	19.4	22.7	25.3	22.8
1999	22.7	35.6	29.5	24.2	21.1	21.4	30.0	21.3
2000	20.1	32.4	30.5	22.6	17.0	20.3	27.7	19.8
2001	na	na	na	na	na	na	na	na
FEMALES								
1991	6.2	6.5	7.9	7.2	8.6	5.2	8.1	5.5
1992	5.6	6.8	6.3	6.3	7.4	7.4	9.1	5.2
1993	3.7	5.4	6.5	7.3	5.0	5.8	6.7	4.4
1994	4.1	6.2	7.2	6.2	6.5	4.6	6.3	4.6
1995	6.4	7.1	8.0	8.9	6.1	5.5	5.4	5.5
1996	4.4	6.8	7.5	6.8	6.2	5.7	5.0	4.9
1997	6.6	7.5	8.4	8.3	6.2	6.9	6.8	5.8
1998	6.3	7.5	8.8	6.0	6.4	5.6	6.6	5.5
1999	5.5	8.1	7.2	7.4	5.5	4.1	3.3	5.1
2000	6.0	7.3	8.8	5.2	5.4	5.9	5.0	5.2
2001	na	na	na	na	na	na	na	na
PERSONS								
1991	15.9	18.3	18.4	17.0	14.0	13.9	19.9	13.2
1992	16.7	17.7	15.5	15.8	15.6	16.4	16.6	12.8
1993	14.0	16.6	13.8	15.3	13.6	13.5	15.8	11.5
1994	15.9	18.2	17.1	16.2	14.8	13.3	15.9	12.7
1995	15.8	20.2	17.9	16.4	14.3	12.2	13.8	13.0
1996	15.8	19.8	18.2	14.6	15.2	14.1	16.5	13.2
1997	18.7	22.1	19.1	16.3	13.9	13.8	17.2	14.2
1998	16.6	23.8	20.5	15.4	12.9	13.8	13.8	14.0
1999	14.3	21.7	18.3	15.9	13.3	12.4	13.6	13.0
2000	13.2	19.8	19.6	13.9	11.2	12.9	13.8	12.3
2001	na	na	na	na	na	na	na	na

(a) Deaths per 100,000 of the estimated mid-year population for each age and sex group.

(b) Standardised death rate per 100,000 persons. Standardised to the estimated mid-year 1991 Australian population.

(c) Data for each year of occurrence includes deaths registered in the particular occurrence year and the year immediately following.

SUICIDES IN THE 15-19 & 20-24 YEAR AGE GROUPS

10. SUICIDE BY AGE AND SEX, Number and Rates - Year of registration

Year of registration	Number		Rate(a)	
	15-19	20-24	15-19	20-24
MALES				
1991	133	243	19.0	34.4
1992	125	253	18.4	34.9
1993	111	234	16.7	32.0
1994	122	252	18.6	34.5
1995	98	252	15.1	34.8
1996	114	237	17.4	33.4
1997	121	295	18.4	42.8
1998	116	248	17.5	36.9
1999	97	212	14.4	32.1
2000	89	174	13.0	26.5
2001	95	181	13.6	27.4
FEMALES				
1991	21	65	3.2	9.4
1992	31	46	4.8	6.5
1993	19	36	3.0	5.1
1994	17	40	2.7	5.6
1995	29	55	4.7	7.8
1996	26	30	4.2	4.4
1997	33	60	5.3	9.0
1998	35	47	5.5	7.2
1999	27	44	4.2	6.9
2000	41	34	6.3	5.3
2001	25	38	3.8	5.9
PERSONS				
1991	154	308	11.3	22.1
1992	156	299	11.8	20.9
1993	130	270	10.0	18.7
1994	139	292	10.9	20.3
1995	127	307	10.0	21.5
1996	140	267	10.9	19.1
1997	154	355	12.0	26.1
1998	151	295	11.7	22.2
1999	124	256	9.4	19.7
2000	130	208	9.7	16.1
2001	120	219	8.8	16.8

(a) Deaths per 100,000 of the estimated mid-year population for each age and sex group.

11. SUICIDE BY AGE AND SEX, Number and Rates - Year of occurrence

Year of Occurrence (b)	Number		Rate(a)	
	15-19	20-24	15-19	20-24
MALES				
1991	120	234	17.2	33.1
1992	133	251	19.6	34.6
1993	101	231	15.2	31.6
1994	126	251	19.3	34.4
1995	97	245	14.9	33.8
1996	114	252	17.4	35.5
1997	123	284	18.7	41.2
1998	110	243	16.6	36.1
1999	94	209	14.0	31.7
2000	91	179	13.3	27.3
2001	na	na	na	na
FEMALES				
1991	23	61	3.5	8.8
1992	31	45	4.8	6.4
1993	17	33	2.7	4.6
1994	18	37	2.9	5.2
1995	28	57	4.5	8.1
1996	25	33	4.0	4.8
1997	33	53	5.3	7.9
1998	36	45	5.7	6.9
1999	26	44	4.1	6.9
2000	43	35	6.6	5.5
2001	na	na	na	na
PERSONS				
1991	143	295	10.5	21.1
1992	164	296	12.4	20.7
1993	118	264	9.1	18.3
1994	144	288	11.3	20.0
1995	125	302	9.8	21.1
1996	139	285	10.9	20.4
1997	156	337	12.2	24.8
1998	146	288	11.3	21.7
1999	120	253	9.1	19.4
2000	134	214	10.0	16.6
2001	na	na	na	na

(a) Deaths per 100,000 of the estimated mid-year population for each age and sex group.

(b) Data for each year of occurrence includes deaths registered in the particular occurrence year and the year immediately following.

APPENDIX

STANDARDISED RATES

Age-standardised rates in this paper were calculated using direct standardisation. Five year age groups (0-4, 5-9,.... 85 & over) were used and a small number of records where age was not stated were excluded from the calculations. The standard population was the 1991 Estimated Resident Population (ERP) for Australia.

ESTIMATED RESIDENT POPULATIONS

Estimated Resident Populations used in the calculation of rates in this paper were obtained from the ABS Information Warehouse in November 2002 and were current at that time. Estimated Resident Populations are subject to revision. Small changes in estimates of resident populations in specific age groups may cause slight changes in age-specific or age-standardised rates calculated using those populations. Thus, there may be slight differences between rates published in this paper and rates for the same time period(s) published elsewhere.

Following the 2001 Census, revised Estimated Resident Populations for 1997-2001 will become available in March/April 2003. Rates calculated using these revised ERPs may also be slightly different from rates published in this paper.

CODING

In this paper, suicide deaths data for registration years 1991-1996 were coded to ICD-9 while data for registration years 1997-2001 were coded to ICD-10. For suicide deaths, ICD-9 and ICD-10 classifications are comparable.

Method of suicide	ICD-9 codes (a)	ICD-10 codes(b)
Poisoning by drugs	E950.0 - E950.5	X60 - X64
Poisoning by "other" (includes motor vehicle exhaust)	E950.6 - E952.9	X65 - X69
Hanging, strangulation, and suffocation	E953	X70
Firearms and explosives	E955	X72 - X75
Other (Includes drowning, cutting and piercing instruments, jumping from high places and other and unspecified means)	E954, E956 - E959	X71, X76 - X84
Suicides (total)	E950 - E959	X60 - X84

(a) ICD-9 codes used for 1991-1996 data in this paper

(b) ICD-10 codes used for 1997-2001 data in this paper

TIME LAG BETWEEN OCCURRENCE AND REGISTRATION

The following table shows the pattern of time lags between year of occurrence and year of registration. Most suicide deaths are registered in the same year in which the death occurred, or the year immediately following. The number of suicide deaths registered after the second year is extremely small (less than 0.2%).

SUICIDES : YEAR OF OCCURRENCE BY YEAR OF REGISTRATION

Year of occurrence	Year of registration											TOTAL(b)
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	
Number of suicide deaths												
1991	2,020	268	2	-	2	-	-	-	-	-	-	2,288
1992	-	2,021	231	3	-	-	-	-	-	-	-	2,252
1993	-	-	1,847	192	3	-	-	-	1	-	-	2,039
1994	-	-	-	2,061	222	-	-	1	1	1	-	2,283
1995	-	-	-	-	2,141	218	4	1	1	-	-	2,359
1996	-	-	-	-	-	2,174	247	5	-	-	-	2,421
1997	-	-	-	-	-	-	2,469	174	3	-	1	2,643
1998	-	-	-	-	-	-	-	2,501	132	5	-	2,633
1999	-	-	-	-	-	-	-	-	2,354	134	1	2,488
2000	-	-	-	-	-	-	-	-	-	2,222	166	2,388
2001	-	-	-	-	-	-	-	-	-	-	2,285	na
TOTAL(a)	2,360	2,294	2,081	2,258	2,368	2,393	2,720	2,683	2,492	2,363	2,454	na

(a) Includes suicide registrations for years of occurrence not shown in table.


(b) For each year of occurrence, the total includes deaths registered in the particular occurrence year and the year immediately following.

na not available

COMPARISON BETWEEN OCCURRENCE AND REGISTRATION

The graph on the following page shows age-standardised rates for suicide for by year of occurrence and by year of registration. It is clear from the graph that there are only slight differences between age-standardised rates calculated by year of registration and those calculated by year of occurrence. In general, both ways of presenting the data lead to similar conclusions.

AGE-STANDARDISED DEATH RATES FOR SUICIDE, year of registration & year of occurrence (b)



(a) per 100,000 estimated resident population. Standardised to the estimated mid-year 1991 Australian population.

(b) Data for each occurrence year includes deaths registered in the particular occurrence year and the year immediately following.

(c) Year of Occurrence data for 2001 not yet available.